

MOTIVIC INVARIANTS OF ARC-SYMMETRIC SETS AND BLOW-NASH EQUIVALENCE

GOULWEN FICHO

ABSTRACT. We define invariants of the blow-*Nash* equivalence of real analytic function germs, in a similar way that the motivic zeta functions of Denef & Loeser [5]. As a key ingredient, we extend the virtual Betti numbers, which were known for real algebraic sets [22], as a generalized Euler characteristics for projective constructible arc-symmetric sets. Actually we prove more: the virtual Betti numbers are not only algebraic invariant, but also *Nash*-invariant of arc-symmetric sets. Our zeta functions enable to sketch the blow-*Nash* equivalence classes of Brieskorn polynomials of two variables.

INTRODUCTION

In the study of real analytic function germs, the choice of a good equivalence relation is an important question. Whereas a topological equivalence is too coarse and a C^1 -equivalence too fine, blow-analytic equivalence, a notion introduced by T.-C. Kuo in 1985 (see [17], and [10] for a survey) seems to behave better, especially with respect to finiteness properties. In this paper we will focus on a particular case of blow-analytic equivalence, say blow-*Nash* equivalence. Let $f, g : (\mathbb{R}^d, 0) \rightarrow (\mathbb{R}, 0)$ be real analytic function germs; f and g are said to be blow-*Nash* equivalent if there exist two algebraic modifications $\pi_f : (M_f, \pi_f^{-1}(0)) \rightarrow (\mathbb{R}^d, 0)$ and $\pi_g : (M_g, \pi_g^{-1}(0)) \rightarrow (\mathbb{R}^d, 0)$, and a *Nash*-isomorphism, that is an analytic isomorphism with semi-algebraic graph, $\phi : (M_f, \pi_f^{-1}(0)) \rightarrow (M_g, \pi_g^{-1}(0))$ which respects the multiplicity of the jacobian determinants of π_f and π_g and which induces a homeomorphism $h : (\mathbb{R}^d, 0) \rightarrow (\mathbb{R}^d, 0)$ such that $f = g \circ h$. Here by a modification of f we mean a proper birational map which is an isomorphism over the complement of the zero locus of f . One can define such an equivalence relation on germs of real analytic varieties, and S. Koike proved ([13, 14]) that finiteness properties hold in this case. However in the case of germs of functions, the question of moduli is still an open question.

A common issue for blow-analytic equivalence and blow-*Nash* equivalence is to prove that, when it is the case, two given function germs are not equivalent. The difficulty rises in the lack of invariants known for these equivalence relations. Up to now, just one kind of invariants have been known: the Fukui invariants. To an analytic function germ f , the Fukui invariants associate the set of possible orders n of series $f \circ \gamma(t) = a_n t^n + \dots$, $a_n \neq 0$, for $\gamma : (\mathbb{R}, 0) \rightarrow (\mathbb{R}^d, 0)$ an analytic arc ([9, 12]). There exists also a version of the Fukui invariants related to the sign of f .

Using motivic integration combined with the construction of a computable motivic invariant for arc-symmetric sets, the virtual Betti numbers, we introduce in this paper zeta functions $Z(T), Z^\pm(T)$ of a real analytic function germ that belong to $\mathbb{Z}[u, u^{-1}][[T]]$, and take into account not only the orders of series $f \circ \gamma(t)$ but also the geometry of the sets $\chi_n(f)$ of arcs γ that realize a given order n (for precise definitions, see section 4.1). These zeta functions are

1991 *Mathematics Subject Classification.* 14B05, 14P20, 14P25, 32S15.

similar to the motivic zeta functions of Denef & Loeser [5]. We prove that actually they are invariants for blow-*Nash* equivalence. The proof is directly inspired by the work of Denef & Loeser via their formulae for the zeta functions in terms of a modification of the zero locus of the analytic function germ (propositions 4.2, 4.4). It uses the powerful machinery of motivic integration, a theory introduced by M. Kontsevitch in 1995 [16] and developed by J. Denef and F. Loeser [5, 6, 7, 8], in particular the fundamental change of variable formula (4.16).

In order to dispose of computable invariants, motivic integration requires computable measures, or in other words Generalized Euler Characteristics. A Generalized Euler Characteristic is an additive and multiplicative invariant defined on the level of the Grothendieck group of varieties. In our setting of the blow-*Nash* equivalence, we need invariants of the Zariski constructible sets over real algebraic varieties $\chi_n(f)$ (real algebraic variety in the sense of [4]), and we ask it to be respected by *Nash*-isomorphisms. It leads naturally to the category of *Nash*-varieties, and more generally of arc-symmetric sets.

Arc-symmetric sets have been introduced in 1988 by K. Kurdyka [18] in order to study “rigid components” of semi-algebraic sets. Arc-symmetric sets enabled him to prove Borel theorem [19] which says that injective endomorphisms of real algebraic sets are surjective. With a slightly different definition of arc-symmetric sets A. Parusiński has proved the same result by using the fact that these sets form a constructible category [23].

In section 2, we give conditions on an invariant defined on connected components of compact nonsingular real algebraic varieties such that it extends to a additive invariant on the constructible category of arc-symmetric sets. Additive means that $\chi(A) = \chi(B) + \chi(A \setminus B)$ for $B \subset A$ an inclusion of arc-symmetric sets. Let us stress the fact that the unique such additive invariant known up to now in the real case is the classical Euler characteristic with compact support, and as a matter of fact it is the unique Generalized Euler Characteristic for semi-algebraic sets up to homeomorphism [24]. As a fundamental example, we prove that the Betti numbers with $\mathbb{Z}_2$ -coefficient defined on connected components A of compact nonsingular real algebraic sets by $b_k(A) = \dim H_k(A, \mathbb{Z}_2)$, give such an invariant β_k on arc-symmetric sets for each $k \in \mathbb{N}$ (corollary 2.4), called k -Virtual Betti number. We make them multiplicative by putting $\beta(A) = \sum_{k=0}^{\dim A} \beta_k(A) u^k \in \mathbb{Z}[u]$, called virtual Poincaré polynomial of A . This invariant is different from the classical Euler characteristic with compact support, and in particular it enables to distinguish homeomorphic real algebraic sets which are not isomorphic. Moreover it respects dimension as put in light by the formula $\deg(\beta(A)) = \dim(A)$ (see remark 2.8) whereas the Euler characteristic with compact support identifies the dimension. These numbers have been proven to be additive invariant of real algebraic varieties recently by C. McCrory and Adam Parusiński in [22]; in this paper we extend the virtual Betti numbers to the more general context of arc-symmetric sets, and we prove the invariance not only under algebraic isomorphisms but also under *Nash*-isomorphisms (see 3.2).

Note that the virtual Betti numbers over real algebraic sets have been introduced independently by C. McCrory and A. Parusiński [15], and by B. Totaro [25]. Moreover S. Koike and A. Parusiński have defined in the same way zeta functions, by using the classical Euler characteristic with compact support, which are invariant for blow-analytic equivalence ([15]). The advantage of our zeta functions, whose invariance is proven only for blow-*Nash* equivalence, is that the Virtual Betti numbers have a better behaviour with respect to algebraicity and analyticity than the classical Euler characteristic with compact support which is merely topological.

In the first section we study the behaviour of arc-symmetric sets with respect to closure, irreducibility, nonsingularity and overall resolution of singularities (proposition 1.12). In the

second one we show that, under certain conditions, an invariant defined over connected components of compact nonsingular algebraic sets can be extended to an additive invariant on all arc-symmetric sets (cf theorem 2.3). Here by invariant we mean up to algebraic isomorphisms (see 1.10). This result requires the use of the Weak Factorization Theorem ([1, 26]) and of the desingularisation Theorem of H. Hironaka [11], and is in the spirit of the description of the naive Grothendieck group of k -varieties, with k a field of characteristic zero, given by F. Bittner in [3]. In particular this part leads to the existence of the Generalized Euler Characteristic β , defined from the Betti numbers, for arc-symmetric sets. Then we extend the invariance of β in section 3, not only to algebraic isomorphisms of arc-symmetric sets but also to *Nash*-isomorphisms (for a precise definition see 3.2). In the final part we define zeta functions of a real analytic function germ and we prove, by the way of motivic integration, the main theorem of this part: $Z(T), Z^\pm(T)$ are invariants for blow-*Nash* equivalence. In particular we state Denef & Loeser formulae for the zeta functions (proposition 4.2, 4.4). Finally we apply these invariants to sketch the blow-*Nash* equivalence classes of Brieskorn polynomials in two variables, and to give some examples in three variables.

Acknowledgement. I am greatly indebted to Adam Parusiński, my thesis advisor, for his help during this work at the university of Angers.

1. ARC-SYMMETRIC SETS

1.1. Arc-Symmetric sets and closure. We fix a compactification of $\mathbb{R}^n$, for instance $\mathbb{R}^n \subset \mathbb{P}^n$.

Definition 1.1. Let $A \subset \mathbb{P}^n$ be a semi-algebraic set. We say that A is arc-symmetric if one of the two equivalent conditions holds:

- (1) for every real analytic arc $\gamma :]-\epsilon, \epsilon[\longrightarrow \mathbb{P}^n$ such that $\gamma(]-\epsilon, 0]) \subset A$ there exists $\epsilon' > 0$ such that $\gamma(]0, \epsilon']) \subset A$,
- (2) for every real analytic arc $\gamma :]-\epsilon, \epsilon[\longrightarrow \mathbb{P}^n$ such that $\text{int } \gamma^{-1}(A) \neq \emptyset$ there exists $x_0, \dots, x_n \in]-\epsilon, \epsilon[$ such that $\gamma(]-\epsilon, \epsilon[\setminus \{x_0, \dots, x_n\}) \subset A$.

Remark that an arc-symmetric set need not to be an analytic variety (cf [18], example 1.2).

This definition is the one of A. Parusiński [23]. Note that a closed arc-symmetric set is necessarily compact. This definition differs from the one of K. Kurdyka [18] who only considers closed arc-symmetric sets in $\mathbb{R}^n$. One can think our arc-symmetric sets as projective constructible arc-symmetric sets. Remark that the arc-symmetric sets form a constructible category $\mathcal{AS}$ of semi-algebraic sets:


- $\mathcal{AS}$ contains the algebraic sets,
- $\mathcal{AS}$ is stable under set-theoretic operations $\cup, \cap, \setminus$,
- $\mathcal{AS}$ is stable by inverse images of $\mathcal{AS}$ -map (i-e whose graph is in $\mathcal{AS}$) and images of injective $\mathcal{AS}$ -map,
- each $A \in \mathcal{AS}$ has a well-defined fundamental class with coefficients in $\mathbb{Z}_2$.

In particular there is a notion of closure in $\mathcal{AS}$ (we refer to [23] for a proof).

Proposition 1.2. *Every $A \in \mathcal{AS}$ admits a smallest arc-symmetric set, denoted by $\overline{A}^{\mathcal{AS}}$, containing A and closed in $\mathbb{P}^n$.*

Remark 1.3. Even if $A \in \mathcal{AS}$, we do not have that $\overline{A} \in \mathcal{AS}$. Consider for A the regular part of the Whitney umbrella $zx^2 = y^2$ (see figure 1). The closure of A in $\mathcal{AS}$ is the entire Whitney umbrella.

FIGURE 1. The Whitney umbrella.



One can consider irreducible arc-symmetric sets: $A \in \mathcal{AS}$ is irreducible if $A = B \cup C$, with B and C closed in A and arc-symmetric, implies that $B \subset C$ or $C \subset B$. Remark that an irreducible arc-symmetric set is not necessarily connected with our definition of arc-symmetric sets (as an example consider a hyperbola in the plane). Nevertheless, as proven in [18], an arc-symmetric set A admits a unique decomposition as a finite union of irreducible arc-symmetric sets closed in A . Note that $\mathcal{AS}$ -closure has a good behaviour with respect to irreducibility:

Proposition 1.4. *If $A \in \mathcal{AS}$ is irreducible, then so is $\overline{A}^{\mathcal{AS}}$.*

Proof. Suppose $\overline{A}^{\mathcal{AS}} = B \cup C$ with B and C arc-symmetric and closed in $\overline{A}^{\mathcal{AS}}$. Then B and C are closed and A splits in $A = A \cap \overline{A}^{\mathcal{AS}} = (A \cap B) \cup (A \cap C)$, with $A \cap B$ and $A \cap C$ arc-symmetric and closed in A . But A is irreducible so, for instance, $A \cap B \subset A \cap C$. Then $A = A \cap C$, so $A \subset C$ and $\overline{A}^{\mathcal{AS}} = C$ because C is arc-symmetric and closed. □

Remark 1.5. The hypothesis “ B and C closed” is essential. Let A be the regular part of the Whitney umbrella. The $\mathcal{AS}$ -closure of A is the entire Whitney umbrella which is the disjoint union of two arc-symmetric sets: A and the vertical line.

Recall that the dimension of an arc-symmetric set is its dimension as a semi-algebraic set, and also the dimension of its Zariski closure in the projective space [18] (here by real algebraic variety we mean in the sense of [4]). In particular, if $A \in \mathcal{AS}$ one has $\dim A = \dim \overline{A}^{\mathcal{AS}} = \dim \overline{A}^{\mathbb{Z}}$.

Proposition 1.6. *Let $A \in \mathcal{AS}$. Then $\overline{A}^{\mathcal{AS}} = A \cup \overline{\overline{A} \setminus A}^{\mathcal{AS}}$. In particular $\dim \overline{A}^{\mathcal{AS}} \setminus A < \dim A$.*

Proof. As a union of arc-symmetric sets $F = A \cup \overline{\overline{A} \setminus A}^{\mathcal{AS}}$ is arc-symmetric. Moreover $F = A \cup (\overline{A} \setminus A) \cup (\overline{\overline{A} \setminus A}^{\mathcal{AS}} \setminus A) = \overline{A} \cup \overline{\overline{A} \setminus A}^{\mathcal{AS}}$ thus F is closed. So we have proved that $\overline{A}^{\mathcal{AS}} \subset F$. Moreover $\overline{A} \subset \overline{A}^{\mathcal{AS}}$ because $\overline{A}^{\mathcal{AS}}$ is closed, thus $\overline{\overline{A} \setminus A}^{\mathcal{AS}} \subset \overline{A}^{\mathcal{AS}}$, hence $F \subset \overline{A}^{\mathcal{AS}}$. Consequently $F = \overline{A}^{\mathcal{AS}}$. □

We can adapt proposition 1.5 of [19] to our definition of arc-symmetric sets.

Proposition 1.7. *Let $A \in \mathcal{AS}$ be irreducible, and $B \subset A$ be a closed arc-symmetric subset of A of the same dimension. Then $B = A$.*

Proof. A can be decomposed into the union of two arc-symmetric sets closed in A : $A = B \cup (\overline{A \setminus B})^{\mathcal{AS}} \cap A$. Then, by irreducibility of A , either $B \subset \overline{A \setminus B}^{\mathcal{AS}} \cap A$ or $\overline{A \setminus B}^{\mathcal{AS}} \cap A \subset B$. In the second case $B = A$, and in the first one $B \subset (\overline{A \setminus B}^{\mathcal{AS}}) \setminus (A \setminus B)$. But this can not happen because the dimension of this arc-symmetric set is strictly less than $\dim B$ by proposition 1.6.

□

1.2. Nonsingular arc-symmetric sets. Let us define a nonsingular arc-symmetric set with relation to its Zariski closure in the projective space.

Definition 1.8. An arc-symmetric set A is nonsingular if $A \cap \text{Sing}(\overline{A}^Z) = \emptyset$.

Lemma 1.9. *A nonsingular and connected arc-symmetric set is irreducible.*

Proof. Let $C \in \mathcal{AS}$ be nonsingular and connected, and $A \subset C$ a closed connected arc-symmetric set of the same dimension. We are going to prove that $A = C$.

Denote by A_0 the semi-algebraic set consisting of the part of maximal dimension of A , and put $A_1 = \overline{A_0}$. Then A_1 is a closed semi-algebraic subset of C , contained in A . But A_1 is also open in C : take $a \in A_1$; there exists an open ball D with center a included in C . Then $\dim D \cap A_0 = \dim A$, therefore $D \subset A$ because A is arc-symmetric (one can fill D with analytic arcs whose interiors intersect A_0 in non empty sets). Moreover $D \subset A_0$, by definition of A_0 , and then A_1 is an open neighbourhood of a . Finally A_1 is a connected component of C , so $A_1 = C$. But $A_1 \subset A$, therefore $A = C$.

□

Let us state the definition of an isomorphism between arc-symmetric sets.

Definition 1.10. Let $A, B \in \mathcal{AS}$. Then A is isomorphic to B if and only if there exist Zariski open subsets U and V in $\overline{A}^Z$ and $\overline{B}^Z$ containing A and B , and an algebraic isomorphism $\phi : U \longrightarrow V$ such that $\phi(A) = B$.

The following proposition says that closed and nonsingular arc-symmetric sets are very similar to compact nonsingular real algebraic varieties.

Proposition 1.11. *Let $A \in \mathcal{AS}$ be compact and nonsingular. Then A is isomorphic to a union of connected components of some compact nonsingular real algebraic variety.*

Proof. Let $X = \overline{A}^Z$ be the Zariski closure of A in the projective space, and $\pi : \tilde{X} \longrightarrow X$ a resolution of singularities of X . Remark that $\dim A = \dim X = \dim \tilde{X}$ and that A is isomorphic to the subset $\pi^{-1}(A) = \tilde{A}$ of $\tilde{X}$ because $A \subset \text{Reg}(X)$ and $\text{Reg}(X)$ is a Zariski open subset of X isomorphic to $\pi^{-1}(\text{Reg}(X)) \subset \tilde{X}$. Denote by $\tilde{X} = \bigcup_{i \in I} C_i$ the decomposition of $\tilde{X}$ in connected components. Each $C_i, i \in I$ is a closed and nonsingular arc-symmetric set, hence irreducible by proposition 1.9. Therefore $\tilde{A} \cap C_i = C_i$ or $\emptyset$ because of proposition 1.7, and $\tilde{A}$ is a union of connected components of $\tilde{X}$ as claimed.

□

1.3. Arc-Symmetric sets and resolution of singularities. The following proposition is just an adaptation of Theorem 2.6 of [18] to our definition of arc-symmetric sets.

Proposition 1.12. *Let $A \in \mathcal{AS}$ be irreducible. Let X be a real algebraic variety containing A with $\dim X = \dim A$, and $\pi : \tilde{X} \rightarrow X$ a resolution of singularities for X (cf [11]). There exists a unique connected component $\tilde{A}$ of $\tilde{X}$ such that $\pi(\tilde{A}) = \overline{\text{Reg}(A)}$.*

Proof. Let $\tilde{A}_0$ be an irreducible arc-symmetric component of dimension $\dim A$ of $\pi^{-1}(A)$ (such an $\tilde{A}_0$ exists because $\dim \pi^{-1}(A) = \dim A$). Then $\tilde{A}_0$ is contained in some connected component $\tilde{A}$ of $\tilde{X}$. Actually $\tilde{A}_0 \subset \overline{\tilde{A}_0}^{\mathcal{AS}}$, which is irreducible because so is $\tilde{A}_0$ (proposition 1.4) and which is closed, is connected (proposition 1.11). Now $\overline{\tilde{A}_0}^{\mathcal{AS}}$ is included in some connected component of $\tilde{X}$, and is equal to it by proposition 1.7. We can put $\tilde{A} = \overline{\tilde{A}_0}^{\mathcal{AS}}$.

Let us show that $\pi(\tilde{A}) = \overline{\text{Reg}(A)}$. In fact, it suffices to show that $\dim \overline{\pi(\tilde{A})}^{\mathcal{AS}} \setminus \pi(\tilde{A}) < \dim A$, what will be done in the next lemma. Indeed, on one hand $\overline{\pi(\tilde{A})}^{\mathcal{AS}} = \overline{A}^{\mathcal{AS}}$ by proposition 1.7, so $\dim \overline{A}^{\mathcal{AS}} \setminus \pi(\tilde{A}) < \dim A$. Now $\text{Reg}(A) \cap (\overline{A}^{\mathcal{AS}} \setminus \pi(\tilde{A}))$ is an open subset of $\overline{A}^{\mathcal{AS}}$ of dimension strictly less than $\dim A$, so $\text{Reg}(A) \cap \overline{A}^{\mathcal{AS}} \setminus \pi(\tilde{A}) = \emptyset$. This implies $\text{Reg}(A) \subset \pi(\tilde{A})$. On the other hand, if E denotes the exceptional divisor of the resolution, one has $\pi(\tilde{A} \setminus E) \subset \text{Reg}(\overline{A}^{\mathcal{AS}})$. However $\text{Reg}(\overline{A}^{\mathcal{AS}}) \subset \overline{\text{Reg}(A)}$ because $\dim \overline{A}^{\mathcal{AS}} \setminus A < \dim A$ by proposition 1.6.

Thus we have the following inclusions $\pi(\tilde{A} \setminus E) \subset \overline{\text{Reg}(A)} \subset \pi(\tilde{A})$ which gives the conclusion by taking the closure. □

Lemma 1.13. *Let A and $\tilde{A}$ be as in the proof of proposition 1.12. Then*

$$\dim \overline{\pi(\tilde{A})}^{\mathcal{AS}} \setminus \pi(\tilde{A}) < \dim A.$$


Proof. Let us show that $\overline{\pi(\tilde{A})}^{\mathcal{AS}} \subset F = \pi(\tilde{A}) \cup \overline{\pi(E)}^{\mathcal{AS}}$. Remark that if F is closed and arc-symmetric the result is proved. As π is proper $\pi(\tilde{A})$ is closed and so is F . Now, let $\gamma :]-\epsilon, \epsilon[\rightarrow \mathbb{P}^n$ be an real analytic arc such that $\text{int } \gamma^{-1}(F) \neq \emptyset$. Then either $\text{int } \gamma^{-1}(\overline{\pi(E)}^{\mathcal{AS}}) \neq \emptyset$ and $\gamma(]-\epsilon, \epsilon[) \subset \overline{\pi(E)}^{\mathcal{AS}}$, or $\text{int } \gamma^{-1}(\pi(\tilde{A}) \setminus \overline{\pi(E)}^{\mathcal{AS}}) \neq \emptyset$. In the latter case there exists a unique analytic arc $\tilde{\gamma} :]-\epsilon, \epsilon[\rightarrow \mathbb{P}^n$ such that $\pi \circ \tilde{\gamma} = \gamma$. One has $\text{int } \tilde{\gamma}^{-1}(\tilde{A}) \neq \emptyset$, so $\tilde{\gamma}(]-\epsilon, \epsilon[) \subset \tilde{A}$ and finally $\gamma(]-\epsilon, \epsilon[) \subset \pi(\tilde{A}) \subset F$. Thus F is arc-symmetric. □

Remark 1.14. Denote by D the singular locus of X and by E the exceptional divisor of the resolution. Then $\pi : \tilde{A} \setminus E \rightarrow \overline{A}^{\mathcal{AS}} \setminus D$ is an isomorphism of arc-symmetric sets (restriction of an algebraic isomorphism).

If we add the assumption that A is nonsingular, then the conclusion becomes $\pi(\tilde{A}) = \overline{A}$. Moreover $\pi : \tilde{A} \setminus (\tilde{A} \setminus \pi^{-1}(A)) \rightarrow A$ is an isomorphism of arc-symmetric sets, and $\dim \tilde{A} \setminus \pi^{-1}(A) < \dim A$.

One can describe with precision the differences between A and $\pi(\tilde{A})$. Actually the symmetric difference of A and $\pi(\tilde{A})$ consists of a semi-algebraic set of dimension strictly less than $\dim A$, and more precisely one has the following proposition.

FIGURE 2. Resolution of the Whitney umbrella.



Proposition 1.15. *Let A and $\tilde{A}$ be as in proposition 1.12. Then*

$$A \setminus (\pi(\tilde{A}) \cap A) = \{x \in \text{Sing}(A), \dim_x A < \dim A\}$$

and

$$\begin{aligned} \pi(\tilde{A}) \setminus (A \cap \pi(\tilde{A})) &= \text{Reg}(\overline{A}^{\text{AS}}) \setminus A \cup \{x \in \text{Sing}(\overline{A}^{\text{AS}} \setminus A), \dim_x \overline{A}^{\text{AS}} = \dim A\} \\ &= \{x \in \overline{A}^{\text{AS}} \setminus A, \dim_x \overline{A}^{\text{AS}} = \dim A\}. \end{aligned}$$

Proof. For the first equality, remark that $A \setminus (\pi(\tilde{A}) \cap A) \subset \text{Sing}(A)$ and that $\text{Sing}(A)$ splits in $\{x \in \text{Sing}(A), \dim_x A < \dim A\} \cup \{x \in \text{Sing}(A), \dim_x A = \dim A\}$. But recall that $\pi(\tilde{A}) = \overline{\text{Reg}(A)}$, and so $\text{Sing}(A) \cap \pi(\tilde{A}) = \text{Sing}(A) \cap \overline{\text{Reg}(A)} = \{x \in \text{Sing}(A), \dim_x A = \dim A\}$.

In the same way, $\pi(\tilde{A}) \setminus (A \cap \pi(\tilde{A})) \subset \overline{A}^{\text{AS}} \setminus A$ and $\overline{A}^{\text{AS}} \setminus A = \text{Reg}(\overline{A}^{\text{AS}} \setminus A) \cup S_1 \cup S_2$ with $S_1 = \{x \in \text{Sing}(\overline{A}^{\text{AS}} \setminus A), \dim_x \overline{A}^{\text{AS}} = \dim A\}$ and $S_2 = \{x \in \text{Sing}(\overline{A}^{\text{AS}} \setminus A), \dim_x \overline{A}^{\text{AS}} < \dim A\}$. One has $S_2 \cap \overline{\text{Reg}(A)} = \emptyset$ and $S_1 \subset \overline{\text{Reg}(A)}$. Moreover the remark 1.14 induces that $\text{Reg}(\overline{A}^{\text{AS}} \setminus A) \subset \overline{A}^{\text{AS}} \setminus D \subset \pi(\tilde{A})$, hence the conclusion. $\square$

Example 1.16. Consider once more the Whitney umbrella $zx^2 = y^2$. One can resolve the singularities by blowing-up along the z -axis. Let $\sigma(u, v, w) = (u, uv, w)$ denote the chart which contains the hole strict transform of the Whitney umbrella; then it has equation $w = v^2$, while the exceptional divisor has equation $u = 0$ (figure 2).

On one hand, if A is just the regular part minus the circle in dotted line, then $A \subset \pi(\tilde{A})$ and $\pi(\tilde{A}) \setminus (A \cap \pi(\tilde{A}))$ consists of the circle and the closed upper part of the vertical line. On the other and if A is the entire Whitney umbrella, then $\pi(\tilde{A}) \subset A$ and $A \setminus (\pi(\tilde{A}) \cap A)$ is the open bottom part of the vertical line.

Let us finish this section by giving the well-known particular case of the “blow-up” of a closed nonsingular arc-symmetric set. We state the proposition in terms of connected component of nonsingular compact algebraic varieties (recall proposition 1.11).

Proposition 1.17. *Let $Y \subset X$ be compact nonsingular algebraic varieties such that $\dim Y < \dim X$, and $A \subset X$ be a connected component of X . Denote by $\pi : \tilde{X} \rightarrow X$ the blow-up of X along Y . Then π is surjective and $\pi^{-1}(A)$ is a connected component of $\tilde{X}$.*

2. GENERALIZED EULER CHARACTERISTIC OF ARC-SYMMETRIC SETS

Definition 2.1. An additive map on $\mathcal{AS}$ with values in an abelian group is a map χ defined on $\mathcal{AS}$ such that

- (1) for A and B arc-symmetric sets with A isomorphic to B , then $\chi(A) = \chi(B)$,
- (2) for $B \subset A$ inclusion of arc-symmetric sets, then $\chi(A) = \chi(B) + \chi(A \setminus B)$.

If moreover χ satisfies $\chi(A \times B) = \chi(A) \times \chi(B)$ for A, B arc-symmetric sets, then we say that χ is a generalized Euler characteristic on $\mathcal{AS}$.

Remark 2.2. The Euler characteristic with compact supports is a generalized Euler characteristic on $\mathcal{AS}$. Actually if we consider just semi-algebraic sets, with isomorphism replaced by homeomorphism, the Euler characteristic with compact support is the unique generalized Euler characteristic (see [24]). However, for complex algebraic varieties, there exist a lot of such generalized Euler characteristic, for example given by mixed Hodge structure (see [5, 21])

The aim of this section is to give sufficiently good conditions on an invariant χ over the closed (i-e compact) and nonsingular arc-symmetric sets such that χ extends to an additive map on $\mathcal{AS}$. We state the theorem in terms of connected components of real algebraic varieties thanks to proposition 1.11. The method is inspired from the one of [3], and the main result is:

Theorem 2.3. *Let χ be a map defined on connected components of compact nonsingular real algebraic varieties with values in an abelian group and such that*

- P1:** $\chi(\emptyset) = 0$,
- P2:** *if A and B are connected components of compact nonsingular real algebraic varieties which are isomorphic as arc-symmetric sets, then $\chi(A) = \chi(B)$,*
- P3:** *with notations and assumptions of proposition 1.17,*

$$\chi(\pi^{-1}(A)) - \chi(\pi^{-1}(A) \cap \pi^{-1}(A \cap Y)) = \chi(A) - \chi(A \cap Y).$$

Then χ extends to an additive map defined on $\mathcal{AS}$.

Before giving the proof of the theorem, let us give some consequences. First this result enables us to give one example of such an additive map by considering the homology with $\mathbb{Z}_2$ coefficients.

Corollary 2.4. *Let β_i be defined on a connected component A of a compact nonsingular algebraic variety by $\beta_i(A) = \dim H_i(A, \mathbb{Z}_2)$, the Betti number mod-2 of A , for $i \in \mathbb{N}$. Then β_i extends to an additive map defined on $\mathcal{AS}$.*

Proof. Property P1 and P2 of theorem 2.3 are clear. Let us prove property P3. With notations of P3, the exact sequences with coefficients in $\mathbb{Z}/2$ of the pairs $(\tilde{A}, \tilde{A} \cap \pi^{-1}(B))$ and (A, B) give the following commutative diagram

$$\begin{array}{ccccccc} \cdots & \longrightarrow & H_{i-1}(\tilde{A}, \tilde{A} \cap \pi^{-1}(B)) & \longrightarrow & H_i(\tilde{A} \cap \pi^{-1}(B)) & \longrightarrow & H_i(\tilde{A}) \longrightarrow \cdots \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & H_{i-1}(A, B) & \longrightarrow & H_i(B) & \longrightarrow & H_i(A) \longrightarrow \cdots \end{array}$$

where the vertical arrows are induced by π . Note that $\pi_* : H_{i-1}(\tilde{A}, \tilde{A} \cap \pi^{-1}(B)) \rightarrow H_{i-1}(A, B)$ is an isomorphism because π is a homeomorphism between $\tilde{A} \setminus \tilde{A} \cap \pi^{-1}(B)$ and $A \setminus B$, and that $\pi_* : H_i(\tilde{A}) \rightarrow H_i(A)$ is surjective because π is of degree 1 and the varieties are $\mathbb{Z}/2$ -oriented. Now it is an easy game to check that the following sequence

$$0 \longrightarrow H_i(\tilde{A} \cap \pi^{-1}(B)) \longrightarrow H_i(B) \oplus H_i(\tilde{A}) \longrightarrow H_i(A) \longrightarrow 0$$

is exact, hence

$$\beta_i(\tilde{A}) - \beta_i(\tilde{A} \cap \pi^{-1}(B)) = \beta_i(A) - \beta_i(B).$$

Then we can apply theorem 2.3.

□

It turns out to be easy to adapt theorem 2.3 in order to obtain not only additive maps but also generalized Euler characteristics.

Theorem 2.5. *Let χ be as in theorem 2.3. Assume moreover that χ takes values in a commutative ring, and that for connected components of compact nonsingular real algebraic varieties A, B the relation $\chi(A \times B) = \chi(A)\chi(B)$ holds. Then the extension of χ on $\mathcal{AS}$ of theorem 2.3 is a generalized Euler characteristic.*

The following corollary is an immediate consequence of Künneth formula.

Corollary 2.6. *Let β be defined on $A \in \mathcal{AS}$ by*

$$\beta(A) = \sum_{i=0}^{\dim A} \beta_i(A)u^i.$$

Then β is a generalized Euler characteristic on $\mathcal{AS}$.

Example 2.7. (1) If $\mathbb{P}^k$ denote the real projective space of dimension k , which is non-singular and compact, then $\beta(\mathbb{P}^k) = 1 + u + \dots + u^k$. Now, compactify $\mathbb{R}$ in $\mathbb{P}^1$ by adding one point at the infinity. By additivity $\beta(\mathbb{R}) = \beta(\mathbb{P}^1) - \beta(\text{point}) = u$, and so $\beta(\mathbb{R}^k) = u^k$.
(2) Let W be the Whitney umbrella, and L the line included in W . Therefore $\beta(W) = \beta(W \setminus L) + \beta(L)$ and $W \setminus L$ is isomorphic, by blowing-up, to the strict transform of W minus a parabola P . So

$$\beta(W \setminus L) = \beta(\mathbb{A}_{\mathbb{R}}^1 \times P) - \beta(P) = (\beta(\mathbb{A}_{\mathbb{R}}^1) - 1)\beta(P) = (u - 1)u.$$

Finally $\beta(W) = u^2$.

(3) The real algebraic varieties of figure 3 (clearly, one can find reduced equations for such curves) are not isomorphic whereas they are homeomorphic. Indeed one can compute $\beta(C_1)$ and $\beta(C_2)$ by considering the resolutions given by blowing-up the singular point of C_1 and C_2 . One finds $\beta(C_1) = u$ and $\beta(C_2) = 2u - 1$. Remark that the Euler characteristic (one can recover by evaluating u at -1 in this example) does not distinguish these two curves.

Remark 2.8. The invariant β has the following property as we will see in the proof; let $A \in \mathcal{AS}$. Then


$$\dim(A) = \deg(\beta(A)).$$

Let us give the proof of theorem 2.3 first, and later of theorem 2.5.

Proof of theorem 2.3. We prove theorem 2.3 by induction on dimension; the rank n inductive hypothesis claims that χ is defined on arc-symmetric sets of dimension less than or equal to n , invariant under isomorphism of arc-symmetric sets, and additive.

For $n = 0$ the arc-symmetric sets are just finite unions of points and the result is clearly true. Assume that the inductive hypothesis is true for $n - 1$. We prove the result at rank n in two steps:

FIGURE 3. resolution of C_1 and C_2



- (1) if χ is an additive map on the nonsingular elements of $\mathcal{AS}$ of dimension less than or equal to n , then χ extends to an additive map on all arc-symmetric sets of dimension less than or equal to n ,
- (2) if χ satisfies property P1, P2 and P3, then χ extends to an additive map on the nonsingular elements of $\mathcal{AS}$ of dimension less than or equal to n .

Step 1 Let $A \in \mathcal{AS}$ of dimension n . There exists a stratification $\overline{A}^Z = \bigcup_{S \in \mathcal{S}} S$ of $\overline{A}^Z$ with nonsingular algebraic strata, that is the union is a disjoint union of locally closed algebraic varieties (note in particular that we do not ask the strata to be connected). Then $S \cap A$, for each $S \in \mathcal{S}$, is a nonsingular arc-symmetric set, and $\chi(S \cap A)$ is defined. Put $\chi(A) = \sum_{S \in \mathcal{S}} \chi(S \cap A)$. One has to check that $\chi(A)$ is well-defined and satisfies the property of an additive function over $\mathcal{AS}$.

We show firstly that $\chi(A) = \sum_{S \in \mathcal{S}} \chi(S \cap A)$ in the case where A is nonsingular, by induction on the number of elements in $\mathcal{S}$. Indeed take $N_0 \in \mathcal{S}$, then $\chi(A) = \chi(A \setminus (A \cap N_0)) + \chi(A \cap N_0)$ and $\chi(A \setminus (A \cap N_0)) = \sum_{S \in \mathcal{S} \setminus \{N_0\}} \chi(S \cap A)$ by induction, hence the result.

Now, if $\mathcal{S}_1$ and $\mathcal{S}_2$ are two stratifications of $\overline{A}^Z$, one can find a common refinement $\mathcal{S}$. The independence in the nonsingular case induces that $\sum_{S \in \mathcal{S}_1} \chi(S \cap A) = \sum_{S \in \mathcal{S}} \chi(S \cap A) = \sum_{S \in \mathcal{S}_2} \chi(S \cap A)$, then χ does not depend on the choice of the stratification.

Let us show that χ is additive; let $B, A \in \mathcal{AS}$ with $B \subset A$ of dimension less than or equal to n . One can choose a stratification $\bigcup_{S \in \mathcal{S}} S$ of $\overline{A}^Z$ such that $\overline{B}^Z$ and $\overline{A \setminus B}^Z$ are union of strata. Then

$$\sum_{S \in \mathcal{S}} \chi(S \cap B) + \sum_{S \in \mathcal{S}} \chi(S \cap (A \setminus B)) = \sum_{S \in \mathcal{S}} (\chi(S \cap B) + \chi(S \cap (A \setminus B)))$$

and $\chi(S \cap (A \setminus B)) + \chi(S \cap B) = \chi(S \cap A)$ because the strata are nonsingular, so χ is additive.

Step 2 The second step constitutes the heart of the work. Define χ over the nonsingular arc-symmetric sets of dimension n in the following way:

D1: if $A = \bigcup_{i \in I} A_i$ denotes the decomposition of A into irreducible components, put $\chi(A) = \sum_{i \in I} \chi(A_i)$,

D2: if $A \in \mathcal{AS}$ is nonsingular and irreducible, then $\chi(A) = \chi(\tilde{A}) - \chi(\tilde{A} \setminus \pi^{-1}(A))$, where $\tilde{A}$ is the connected component of a resolution of singularities of $\overline{A}^Z$ given by remark 1.14 and E the exceptionnal divisor of the resolution.

We have to prove that χ is well-defined, invariant under isomorphisms and additive over the nonsingular elements of $\mathcal{AS}$.

The following lemma will be usefull in the sequel.

Lemma 2.9. *Let $A, B \in \mathcal{AS}$ be nonsingular, irreducible and isomorphic. Suppose that $\overline{A}^Z$ and $\overline{B}^Z$ are nonsingular, and denote by $\tilde{A} \subset \overline{A}^Z$ and $\tilde{B} \subset \overline{B}^Z$ the connected components containing A and B . Then $\chi(\tilde{A}) - \chi(\tilde{A} \setminus A) = \chi(\tilde{B}) - \chi(\tilde{B} \setminus B)$.*

Proof. By definition of an isomorphism between arc-symmetric sets, we know that $\overline{A}^Z$ and $\overline{B}^Z$ are birationally equivalent, and the weak factorization theorem [1, 26] factors this birational isomorphism in a succession of blow-ups and blow-downs. In particular we can assume that the birational isomorphism is just a blow-up $\pi : \overline{A}^Z \longrightarrow \overline{B}^Z$ along a nonsingular variety C such that $C \cap B = \emptyset$. Note that $\pi^{-1}(\tilde{B}) = \tilde{A}$ by proposition 1.17.

Now $\chi(\tilde{B} \setminus B) = \chi(\tilde{B} \cap C) + \chi(\tilde{B} \setminus (B \cup C))$ by the additivity inductive hypothesis for $\dim \tilde{B} \setminus B < \dim B$ by proposition 1.6. Moreover $\chi(\tilde{B}) - \chi(\tilde{B} \cap C) = \chi(\tilde{A}) - \chi(\tilde{A} \cap \pi^{-1}(C))$ by property P3, and $\chi(\tilde{B} \setminus (B \cup C)) = \chi(\tilde{A} \setminus (A \cup \pi^{-1}(C)))$ by the inductive hypothesis on invariance under isomorphism. Therefore $\chi(\tilde{B}) - \chi(\tilde{B} \setminus B) = \chi(\tilde{A}) - \chi(\tilde{A} \cap \pi^{-1}(C)) - \chi(\tilde{A} \setminus (A \cup \pi^{-1}(C)))$ which is equal to $\chi(\tilde{A}) - \chi(\tilde{A} \setminus A)$ by the additivity inductive hypothesis. $\square$

Let us check that the definition of χ for the nonsingular and irreducible arc-symmetric sets of dimension n does not depend on the choice of the resolution of singularities of remark 1.14.

Let $A \in \mathcal{AS}$ be nonsingular and irreducible, and let $\pi_i : \tilde{X}_i \longrightarrow \overline{A}^Z$ for $i \in \{1, 2\}$ be resolutions of singularities of $\overline{A}^Z$. Let $\tilde{A}_i$ be the connected components of $\tilde{X}_i$ given by proposition 1.12. One has to show that

$$\chi(\tilde{A}_1) - \chi(\tilde{A}_1 \setminus \pi_1^{-1}(A)) = \chi(\tilde{A}_2) - \chi(\tilde{A}_2 \setminus \pi_2^{-1}(A)).$$

But $\pi_1^{-1}(A)$ and $\pi_2^{-1}(A)$ are isomorphic irreducible nonsingular arc-symmetric sets because π_i is an isomorphism on a Zariski open subsets of $\tilde{X}_i$ containing $\pi_i^{-1}(A)$, for $i \in \{1, 2\}$. Therefore lemma 2.9 applies and χ is well-defined.

Now let us show that χ is invariant under isomorphisms of arc-symmetric sets. The proof is very similar to the last one. Let $A, B \in \mathcal{AS}$ be irreducible and isomorphic, which means that there exists Zariski open subsets U and V in $\overline{A}^Z$ and $\overline{B}^Z$, and an algebraic isomorphism $\phi : U \longrightarrow V$ such that $\phi(A) = B$. Choose resolutions of singularities $\pi_A : \tilde{X} \longrightarrow \overline{A}^Z$ and $\pi_B : \tilde{Y} \longrightarrow \overline{B}^Z$ for $\overline{A}^Z$ and $\overline{B}^Z$. Then $\pi_A^{-1}(A)$ and $\pi_B^{-1}(B)$ are isomorphic as arc-symmetric sets, and then by lemma 2.9 $\chi(\pi_A^{-1}(A)) = \chi(\pi_B^{-1}(B))$. Moreover one has $\chi(\pi_A^{-1}(A)) = \chi(A)$ because both are equal to $\chi(\tilde{A}) - \chi(\tilde{A} \setminus \pi_A^{-1}(A))$, where $\tilde{A}$ is the connected component of $\tilde{X}$ given by proposition 1.12. In the same way one has $\chi(\pi_B^{-1}(B)) = \chi(B)$, hence

$$\chi(A) = \chi(\pi_A^{-1}(A)) = \chi(\pi_B^{-1}(B)) = \chi(B).$$

In the case where A and B are not irreducible, it suffices to decompose A and B in irreducible components, and apply the property D1 because an isomorphism between arc-symmetric sets respects the irreducible components.

Finally, let us check that χ is additive. Let $B \subset A$ be an inclusion of nonsingular arc-symmetric sets. Note that by definition of χ we need to prove the result only in the case where A is irreducible.

If $\dim B = \dim A$, then $\overline{B}^{\text{AS}} = \overline{A}^{\text{AS}}$ by proposition 1.7, and $\overline{B}^{\mathcal{Z}} = \overline{A}^{\mathcal{Z}}$. Now choose a resolution of singularities $\pi : \tilde{X} \longrightarrow \overline{A}^{\mathcal{Z}}$ for $\overline{A}^{\mathcal{Z}}$. If $\tilde{A}$ denotes the connected component of $\tilde{X}$ given by proposition 1.12 for A , then it is also the component associated to B , and therefore $\chi(B) = \chi(\tilde{A}) - \chi(\tilde{A} \setminus \pi^{-1}(B))$. Now $\chi(\tilde{A} \setminus \pi^{-1}(B)) = \chi(\tilde{A} \setminus \pi^{-1}(A)) + \chi(\pi^{-1}(A) \setminus \pi^{-1}(B))$ by the inductive hypothesis on additivity. So $\chi(B) = \chi(A) - \chi(A \setminus B)$ because $\chi(\pi^{-1}(A) \setminus \pi^{-1}(B)) = \chi(A \setminus B)$ by the invariance under isomorphism in dimension smaller than n .

If $\dim B < \dim A$, choose a resolution of singularities $\pi : \tilde{X} \longrightarrow \overline{A}^{\mathcal{Z}}$ for $\overline{A}^{\mathcal{Z}}$. Then it is also a resolution of singularities of $\overline{A \setminus B}^{\mathcal{Z}} = \overline{A}^{\mathcal{Z}}$. Then $\chi(A \setminus B) = \chi(\tilde{A}) - \chi(\tilde{A} \setminus \pi^{-1}(A \setminus B))$. Now

$$\chi(\tilde{A} \setminus \pi^{-1}(A \setminus B)) = \chi\left((\tilde{A} \setminus \pi^{-1}(A)) \cup \pi^{-1}(B)\right) = \chi(\tilde{A} \setminus \pi^{-1}(A)) + \chi(\pi^{-1}(B))$$

by the inductive assumption, and once more by the inductive assumption one has $\chi(\pi^{-1}(B)) = \chi(B)$. Finally $\chi(A \setminus B) = \chi(A) - \chi(B)$.

This achieves the proof of step 2, and thus the proof of theorem 2.3. □

As it was the case for the previous proof, we are going to prove theorem 2.5 by induction on dimension. The following relations will be useful:

$$(1) \quad \chi(\sqcup_{i=1}^k A_i) = \sum_{i=1}^k \chi(A_i),$$

where the union of the arc-symmetric sets $A_i, i = 1, \dots, k$ is disjoint, and

$$(2) \quad \chi(A) = \chi(\tilde{A}) - \chi(\tilde{A} \setminus A),$$

where A is a nonsingular arc-symmetric set whose arc-symmetric closure $\tilde{A}$ is nonsingular.

Proof of theorem 2.5. Put, as an inductive hypothesis at rank n , that χ is multiplicative for all arc-symmetric sets of dimension strictly less than or equal to n .

Remark that we can restrict our attention to the nonsingular case because by considering stratifications of arc-symmetric sets with nonsingular strata, we prove the multiplicativity directly with formula (1).

Assume therefore that A, B are nonsingular arc-symmetric sets of dimension less than or equal to n ; suppose that $\dim A = n$ for instance.

In case A compact, the result follows from a finite induction on the dimension of B : indeed, resolving the singularities of $\overline{B}^{\mathcal{Z}}$, one can assume that $B \subset \tilde{B}$, with $\tilde{B}$ the nonsingular arc-symmetric closure of B . Then, by (2)

$$\chi(A \times B) = \chi(A \times \tilde{B}) - \chi(A \times (\tilde{B} \setminus B))$$

However $\chi(A \times \tilde{B}) = \chi(A)\chi(\tilde{B})$ for they are compact and nonsingular, and

$$\chi(A \times (\tilde{B} \setminus B)) = \chi(A)\chi(\tilde{B} \setminus B)$$

as we can see by stratifying $\tilde{B} \setminus B$ with nonsingular strata and using the inductive assumption of the finite induction because $\dim \tilde{B} \setminus B < \dim B$ by lemma 1.6. Consequently

$$\chi(A \times B) = \chi(A)(\chi(\tilde{B}) - \chi(\tilde{B} \setminus B)) = \chi(A)\chi(B).$$

If A is no longer compact, then compactify A and B in $\tilde{A}, \tilde{B}$, and assume that $\tilde{A}, \tilde{B}$ are nonsingular even if it means resolving singularities, as before.

Then, by additivity,

$$\chi(A \times B) = \chi(\tilde{A} \times B) - \chi((\tilde{A} \setminus A) \times \tilde{B}) + \chi((\tilde{A} \setminus A) \times (\tilde{B} \setminus B)).$$

The multiplicativity of the first two terms comes from the preceding case (in the second one, stratify the possibly singular set $\tilde{A} \setminus A$), and the multiplicativity of the third is obtained by the inductive assumption for $\max(\dim \tilde{A} \setminus A, \dim \tilde{B} \setminus B) < n$ by lemma 1.6. Therefore

$$\chi(A \times B) = \chi(\tilde{A})\chi(B) - \chi(\tilde{A} \setminus A)\chi(\tilde{B}) + \chi(\tilde{A} \setminus A)\chi(\tilde{B} \setminus B) = \chi(A)\chi(B),$$

and theorem 2.5 is proven. □

3. INVARIANCE OF β UNDER *Nash*-ISOMORPHISMS

The definition of isomorphism between arc-symmetric sets is algebraic, via birational morphisms. But arc-symmetric sets are also closely related to analytic objects. As an example, the following proposition emphasizes the good behaviour of β with respect to compact algebraic varieties which are nonsingular as analytic varieties. Recall that by $b_i(X)$ we denote the i -th Betti number of X with mod-2 coefficients, and let us put $b(X) = \sum_{i=0}^{\dim X} b_i(X)u^i$.

Proposition 3.1. *Let X be a compact algebraic variety which is nonsingular as an analytic space. Then $\beta(X) = b(X)$.*

Proof. One can desingularize the algebraic singularities of X by a sequence of blowings-up with smooth centers ([2, 11]). At each step of the desingularization one has the following relation, where $Bl_C X$ designs the blowing-up of X along the nonsingular subvariety C with exceptional divisor E :

$$\beta(Bl_C X) - \beta(E) = \beta(X) - \beta(C)$$

because the blowing-up is birational and

$$b(Bl_C X) - b(E) = b(X) - b(C)$$

because X and C are smooth and the blowing-up is a degree one morphism. Remark that $\beta(E) = b(E)$ and $\beta(C) = b(C)$ because E and C are nonsingular and compact arc-symmetric sets, and the same is true for $\tilde{X}$, the desingularization of X . Then $\beta(X)$ and $b(X)$ can be expressed by the same formulae in terms of β for the former and b for the latter, where the spaces involved are nonsingular and compact. Therefore for each one of these spaces β and b coincide, and then $\beta(X) = b(X)$. □

Actually we will see later that the assumption “ X is an algebraic variety” can be replaced by the weaker “ X is a semi-algebraic set”, and so “ X is a *Nash*-manifold”.

In order to relate the analytic aspect of arc-symmetric sets and the behaviour of β , we propose the following definition of *Nash*-isomorphism between arc-symmetric sets.

Definition 3.2. Let $A, B \in \mathcal{AS}$, and assume that there exist compact analytic varieties V_1, V_2 containing A, B respectively, and an analytic isomorphism $\phi : V_1 \longrightarrow V_2$ such that $\phi(A) = B$. If moreover one can choose V_1, V_2 to be semi-algebraic sets and ϕ a semi-algebraic map, then one says that A and B are *Nash-isomorphic*.

One wants this new definition of isomorphism respects the invariance of β and, actually, one has the following result.

Theorem 3.3. *Let $A_1, A_2 \in \mathcal{AS}$ be Nash-isomorphic arc-symmetric sets. Then $\beta(A_1) = \beta(A_2)$.*

Proof. Once more, we are going to prove the result by induction on dimension. As a first step, let us generalize the result of proposition 3.1.

Step 1 Let A be a compact arc-symmetric set which is also a nonsingular analytic subspace of the Zariski closure X of A . Then $\beta(A) = b(A)$.

In order to prove this claim, one wants to apply the same method as in the proof of proposition 3.1, but if C is a smooth center of blowing-up for X it is not true in general that $C \cap A$ is nonsingular, so the equality $\beta(C \cap A) = b(C \cap A)$ does no longer hold. In order to solve this problem, consider the algebraic normalization $\tilde{X}$ of X . There exists $\tilde{A} \subset \tilde{X}$ the analytic normalization of A ([20]), which is analytically isomorphic to A because A is nonsingular as an analytic space. Then $b(A) = b(\tilde{A})$ because b is invariant under homeomorphism.

Moreover $\beta(A) = \beta(\tilde{A})$; actually the algebraic normalization is a birational map, hence an algebraic isomorphism outside compact subvarieties E and D of $\tilde{X}$ and X respectively, of dimension strictly less than $\dim X = \dim A$. Thus $\beta(\tilde{A} \setminus E) = \beta(A \setminus D)$ by corollary 2.4, and the algebraic normalization restricted to $\tilde{A} \cap E$ is an analytic isomorphism onto $A \cap D$, so $\beta(\tilde{A} \cap E) = \beta(A \cap D)$ by the inductive assumption.

Note that $\tilde{X}$ is locally analytically irreducible as a normal space, therefore $\tilde{A}$ is a union of connected components of $\tilde{X}$. Now it is true that $C \cap \tilde{A}$ is nonsingular when C is nonsingular, and the method of the proof of proposition 3.1 applies, therefore $\beta(\tilde{A}) = b(\tilde{A})$. It follows that $\beta(A) = \beta(\tilde{A}) = b(\tilde{A}) = b(A)$.

Step 1 is achieved.

Step 2 Let us prove the theorem in the particular case where A_i is a nonsingular arc-symmetric set for $i = 1, 2$, and moreover, with assumptions of the definition of a *Nash-isomorphism*, the compact analytic variety V_i is supposed to be smooth as an analytic space, for $i = 1, 2$.

- First we show that $\beta(\overline{A_2}^{\mathcal{AS}}) = \beta(\overline{A_1}^{\mathcal{AS}})$.

Remark that $\overline{A_2}^{\mathcal{AS}}$ is a union of connected components of V_2 by proposition 1.12. Thus $\overline{A_2}^{\mathcal{AS}}$ is also nonsingular as an analytic variety and $\beta(\overline{A_2}^{\mathcal{AS}}) = b(\overline{A_2}^{\mathcal{AS}})$ by step 1.

Moreover $\overline{A_2}^{\mathcal{AS}}$ is isomorphic to $\overline{A_1}^{\mathcal{AS}}$ by ϕ : indeed $\phi^{-1}(\overline{A_2}^{\mathcal{AS}})$ is a closed arc-symmetric set because ϕ has an arc-symmetric graph and is continuous, and it contains A_1 so $\overline{A_1}^{\mathcal{AS}} \subset \phi^{-1}(\overline{A_2}^{\mathcal{AS}})$. The reverse inclusion comes from the fact that the image by an injective algebraic map of an arc-symmetric set is still an arc-symmetric set (recall that $\mathcal{AS}$ form a constructible category, cf section 1). Consequently $\overline{A_1}^{\mathcal{AS}}$ is nonsingular as an analytic variety because so is $\overline{A_2}^{\mathcal{AS}}$ and ϕ is an analytic isomorphism, hence $\beta(\overline{A_1}^{\mathcal{AS}}) = b(\overline{A_1}^{\mathcal{AS}})$ by the first step.

Remark also that $b(\overline{A_2}^{\mathcal{AS}}) = b(\overline{A_1}^{\mathcal{AS}})$ because ϕ is a homeomorphism between these two smooth compact topological varieties.

These equalities imply that $\beta(\overline{A_2}^{\mathcal{AS}}) = \beta(\overline{A_1}^{\mathcal{AS}})$.

- Then, remark that $\beta(\overline{A_1}^{\mathcal{AS}} \setminus A_1) = \beta(\overline{A_2}^{\mathcal{AS}} \setminus A_2)$. Indeed this follows from the inductive hypothesis for $\overline{A_1}^{\mathcal{AS}} \setminus A_1$ and $\overline{A_2}^{\mathcal{AS}} \setminus A_2$ are *Nash*-isomorphic arc-symmetric sets of dimension strictly less than $\dim A_2$.
- Finally $\beta(A_1) = \beta(A_2)$. Actually

$$\beta(A_1) = \beta(\overline{A_1}^{\mathcal{AS}}) - \beta(\overline{A_1}^{\mathcal{AS}} \setminus A_1)$$

and

$$\beta(A_2) = \beta(\overline{A_2}^{\mathcal{AS}}) - \beta(\overline{A_2}^{\mathcal{AS}} \setminus A_2),$$

and we have proved that the second members are equal, so $\beta(A_1) = \beta(A_2)$ by additivity of β .

Step 3 Reduction of the problem to Step 2.

By definition of a *Nash*-isomorphism, there exist compact analytic varieties V_1, V_2 containing A, B , and an analytic isomorphism $\phi : V_1 \rightarrow V_2$ such that $\phi(A_1) = A_2$, and moreover V_1 and V_2 are semi-algebraic sets and ϕ is a semi-algebraic map.

Denote by X_1 and X_2 the Zariski closures of V_1 and V_2 .

As a first step, we are going to obtain a regular morphism rather than a semi-algebraic map. Denote by Γ the graph of ϕ . This graph is semi-algebraic and analytic, thus arc-symmetric. Then the projection p_i from $Z = \overline{\Gamma}^Z$ onto X_i , for $i \in \{1, 2\}$, is a regular morphism whose restriction to Γ is an analytic isomorphism with V_i . Moreover the preimages by these restrictions of A_1 and A_2 coincide, so one can put $B = p^{-1}(A)$, where $A = A_i \subset V_i = V \subset X_i = X$ for $i \in \{1, 2\}$ and $p : Z \rightarrow X$ denote the natural projection. Therefore B is an arc-symmetric set *Nash*-isomorphic to A , and the issue is now to prove that $\beta(B) = \beta(A)$.

In order to do this, we want to come down to nonsingular objects.

Desingularize X by a sequence of blowings-up with respect to coherent algebraic sheaves of ideals (this is possible by [2, 11]). Blowing-up Z with respect to the corresponding inverse image ideal sheaves, one has at each step a regular morphism which lifts the projection $p : Z \rightarrow X$ to the corresponding blowing-up by the universal property of algebraic blowing-up. Let $\pi_X : \tilde{X} \rightarrow X$ denote the resolution of singularities of X and $\pi_Z : \tilde{Z} \rightarrow Z$ the corresponding composition of blowings-up of Z . If $\tilde{p}$ denotes the morphism obtained between $\tilde{Z}$ and $\tilde{X}$ by universal property, one has the following commutative diagram:

$$\begin{array}{ccc} \tilde{Z} & \xrightarrow{\tilde{p}} & \tilde{X} \\ \pi_Z \downarrow & & \downarrow \pi_X \\ Z & \xrightarrow{p} & X \end{array}$$

Moreover $\tilde{p}$ restricted to the analytic strict transform $\tilde{\Gamma}$ of Γ is an analytic isomorphism onto the strict transform $\tilde{V}$ of V because so is p between Γ and V (here we consider the blowing-up as an analytic one).

$$\begin{array}{ccc}
\tilde{\Gamma} & \xrightarrow[\sim]{\tilde{p}_{|\tilde{\Gamma}}} & \tilde{V} \\
\pi_{Z|\Gamma} \downarrow & & \downarrow \pi_{X|V} \\
\Gamma & \xrightarrow[\sim]{p_{|\Gamma}} & V
\end{array}$$

Now we reduce to the case where A and B are nonsingular by the inductive hypothesis. Actually the singular part of A and B are not necessarily exchanged by $p_{|\Gamma}$, but $\text{Sing}(A) \cup p_{|\Gamma}^{-1}(\text{Sing}(B))$ and $\text{Sing}(B) \cup p_{|\Gamma}(\text{Sing}(A))$ are *Nash*-isomorphic by the restriction of $p_{|\Gamma}$. Moreover the dimension of these arc-symmetric sets is stricly less than $\dim A = \dim B$, so they have the same image by β by the inductive hypothesis. Let us denote by A' and B' the complement of these sets in A and B . So A' and B' are nonsingular.

As A' is nonsingular, it is isomorphic, in the sense of birational morphism, with its preimage in the desingularization $\tilde{X}$ of X . Consequently B' is also isomorphic to its preimage in $\tilde{\Gamma}$ by commutativity of the first diagram. As a consequence $\beta(A') = \beta(\pi^{-1}(A'))$ and $\beta(B') = \beta(\pi^{-1}(B'))$, and we have reduced the problem to step 2. $\square$

4. ZETA FUNCTIONS AND BLOW-NASH EQUIVALENCE

4.1. Zeta functions. This section is directly inspired by the work of J. Denef & F. Loeser on their motivic zeta functions [5]. We first define zeta functions for an analytic germ of functions. Then we will give a formula to compute these zeta functions in terms of a modification.

Denote by $\mathcal{L}$ the space of analytic arcs at the origin $0 \in \mathbb{R}^d$:

$$\mathcal{L} = \mathcal{L}(\mathbb{R}^d, 0) = \{\gamma : (\mathbb{R}, 0) \longrightarrow (\mathbb{R}^d, 0) | \gamma \text{ is analytic}\},$$

and by $\mathcal{L}_n$ the space of truncated analytic arcs

$$\mathcal{L}_n = \mathcal{L}_n(\mathbb{R}^d, 0) = \{\gamma \in \mathcal{L} | \gamma(t) = a_1 t + a_2 t^2 + \cdots a_n t^n, a_i \in \mathbb{R}^d\}.$$

Let $\pi_n : \mathcal{L} \longrightarrow \mathcal{L}_n$ and $\pi_{n,i} : \mathcal{L}_n \longrightarrow \mathcal{L}_i$, with $n \geq i$, be the truncation morphisms.

Consider $f : (\mathbb{R}^d, 0) \longrightarrow (\mathbb{R}, 0)$ an analytic function germ. We define the naive zeta function $Z_f(T)$ of f as the following element of $\mathbb{Z}[u, u^{-1}]((T))$:

$$Z_f(T) = \sum_{n \geq 1} \beta(\chi_n) u^{-nd} T^n,$$

where

$$\chi_n = \{\gamma \in \mathcal{L}_n | \text{ord}(f \circ \gamma) = n\} = \{\gamma \in \mathcal{L}_n | f \circ \gamma(t) = bt^n + \cdots, b \neq 0\},$$

and the “monodromic” zeta functions, or zeta functions with signs, as

$$Z_f^\pm(T) = \sum_{n \geq 1} \beta(\chi_n^\pm) u^{-nd} T^n,$$

where

$$\chi_n^\pm = \{\gamma \in \mathcal{L}_n | f \circ \gamma(t) = \pm t^n + \cdots\}.$$

Remark that $\chi_n, \chi_n^\pm$, for $n \geq 1$, are constructible subsets of $\mathbb{R}^{nd}$, hence belong to $\mathcal{AS}$.

Example 4.1. Let $f : (\mathbb{R}, 0) \longrightarrow (\mathbb{R}, 0)$ be defined by $f(x) = x^k, k \geq 1$. Then

$$\chi_n = \begin{cases} \{\gamma = a_m t^m + \cdots + a_n t^n | a_m \neq 0\} \simeq \mathbb{R}^* \times \mathbb{R}^{n-m} & \text{if } n=mk \\ \emptyset & \text{otherwise} \end{cases}$$

Therefore $\beta(\chi_n) = (u-1)u^{n-m}$ if $n = mk$ and 0 otherwise, hence

$$Z_f(T) = \sum_{m \geq 1} (u-1)u^{mk-m} \left(\frac{T}{u}\right)^{mk} = (u-1) \frac{T^k}{u - T^k}.$$

To compute the zeta functions with signs, we have to consider the case $k = 2p$ and $k = 2p+1$.

If $k = 2p$, then $\chi_n^- = \emptyset$ and

$$\chi_n^+ = \begin{cases} \{\gamma = \pm t^m + \cdots + a_n t^n | a_m \neq 0\} \simeq \{\pm 1\} \times \mathbb{R}^{n-m} & \text{if } n=mk \\ \emptyset & \text{otherwise,} \end{cases}$$

so

$$Z_f^+(T) = \sum_{m \geq 1} 2u^{mk-m} \left(\frac{T}{u}\right)^{mk} = 2 \frac{T^k}{u - T^k}.$$

If $k = 2p+1$, then

$$\chi_n^\pm = \begin{cases} \{\gamma = \pm t^m + \cdots + a_n t^n | a_m \neq 0\} \simeq \{\pm 1\} \times \mathbb{R}^{n-m} & \text{if } n=mk \\ \emptyset & \text{otherwise,} \end{cases}$$

and

$$Z_f^+(T) = Z_f^-(T) = \sum_{m \geq 1} u^{mk-m} \left(\frac{T}{u}\right)^{mk} = \frac{T^k}{u - T^k}.$$

It may be convenient to express zeta functions in terms of a modification of f , that is a proper birational map which is an isomorphism over the complement of the zero locus of f . Actually there exists a formula, called Denef & Loeser formula, which enables to do this. In the naive case, the Denef & Loeser formula is given by the following proposition.

Proposition 4.2. *Let $\sigma : (M, \sigma^{-1}(0)) \longrightarrow (\mathbb{R}^d, 0)$ be a modification of $\mathbb{R}^d$ such that $f \circ \sigma$ and the jacobian determinant $\text{jac } \sigma$ are normal crossings simultaneously, and assume moreover that σ is an isomorphism over the complement of the zero locus of f .*

Let $(f \circ \sigma)^{-1}(0) = \cup_{j \in J} E_j$ be the decomposition into irreducible components of $(f \circ \sigma)^{-1}(0)$, and assume that $\sigma^{-1}(0) = \cup_{k \in K} E_k$ for some $K \subset J$.

Put $N_i = \text{mult}_{E_i} f \circ \sigma$ and $\nu_i = 1 + \text{mult}_{E_i} \text{jac } \sigma$, and for $I \subset J$ denote by E_I^0 the set $(\cap_{i \in I} E_i) \setminus (\cup_{j \in J \setminus I} E_j)$. Then

$$Z_f(T) = \sum_{I \neq \emptyset} (u-1)^{|I|} \beta(E_I^0 \cap \sigma^{-1}(0)) \prod_{i \in I} \frac{u^{-\nu_i} T^{N_i}}{1 - u^{-\nu_i} T^{N_i}}.$$

Example 4.3. Let $f_k : (\mathbb{R}^2, 0) \longrightarrow (\mathbb{R}, 0)$ be defined by $f_k(x, y) = x^k + y^k, k \geq 2$. The blowing-up at the origin give a suitable modification σ for f . Here $(f \circ \sigma)^{-1}(0)$ consists of just the exceptional divisor $\mathbb{P}^1$ in case k even, and furthermore, in case k odd, of the strict transform of f which is a smooth curve crossing transversally the exceptional divisor. Then

$$Z_{f_k} = \begin{cases} (u^2 - 1)u^{-2} \frac{T^k}{1 - u^{-2} T^k} & \text{if } k \text{ is even} \\ (u-1) \frac{u^{-2} T^k}{1 - u^{-2} T^k} (u + (u-1) \frac{u^{-1} T}{1 - u^{-1} T}) & \text{if } k \text{ is odd} \end{cases}$$

Note in particular that for $k \neq k'$ the zeta function Z_{f_k} and $Z_{f'_k}$ are different.

When we are dealing with signs, one defines coverings $\widetilde{E_I^{0,\pm}}$ of E_I^0 in the following way. Let U be an affine open subset of M such that $f \circ \sigma = u \prod_{i \in I} y_i^{N_i}$ on U , with u a unit. Let us put $R_U^\pm = \{(x, t) \in (E_I^0 \cap U) \times \mathbb{R} : t^m = \pm \frac{1}{u(x)}\}$, where $m = \gcd(N_i)$. Then the $R_U^\pm$ glue together along the $E_I^0 \cap U$ to give $\widetilde{E_I^{0,\pm}}$.

Proposition 4.4. *With assumptions and notations of proposition 4.2, then*

$$Z_f^\pm(T) = \sum_{I \neq \emptyset} (u-1)^{|I|-1} \beta(\widetilde{E_I^{0,\pm}} \cap \sigma^{-1}(0)) \prod_{i \in I} \frac{u^{-\nu_i} T^{N_i}}{1 - u^{-\nu_i} T^{N_i}}.$$

We will prove these results in section 4.4.

Example 4.5. (1) Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be defined by $f(x) = u(x) \prod_{i=1}^k x_i^{N_i}$, with $N_i \in \mathbb{N}$. Then f has already normal crossings, and

$$Z_f(T) = (u-1)^k \prod_{i=1}^k \frac{u^{-1} T^{N_i}}{1 - u^{-1} T^{N_i}}.$$

Now, if there exists at least one N_i odd, then $Z_f^+(T) = Z_f^-(T) = \frac{1}{u-1} Z_f(T)$. On the other hand, if all N_i are even, then $Z_f^-(T) = 0$ and $Z_f^+(T) = \frac{2}{u-1} Z_f(T)$ if u is positive, the converse otherwise..

- (2) Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by $f(x, y) = x^2 + y^2$. As f is positive, $Z_f^-(T) = 0$. We obtain a modification by the same way as in example 4.3, and $\widetilde{E_I^{0,+}}$ is here the boundary of a Mobius band, hence homeomorphic to $\mathbb{P}^1$. Therefore $Z_f^+(T) = (u+1) \frac{u^{-2} T^2}{1 - u^{-2} T^2} = \frac{1}{u-1} Z_f(T)$.
- (3) Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by $f(x, y) = x^2 + y^4$. One can solve the singularities of f by two successive blowings-up, and one obtains that the exceptional divisor E has two irreducible components E_1, E_2 with $N_1 = 2, \nu_1 = 2, N_2 = 4, \nu_2 = 3$. Therefore

$$Z_f(T) = (u-1)^2 \frac{u^{-2} T^2}{1 - u^{-2} T^2} \frac{u^{-3} T^4}{1 - u^{-3} T^4} + (u-1) u \frac{u^{-2} T^2}{1 - u^{-2} T^2} + (u-1) u \frac{u^{-3} T^4}{1 - u^{-3} T^4}.$$

Moreover in this case $\widetilde{E_{\{1\}}^{0,+}}$ and $\widetilde{E_{\{2\}}^{0,+}}$ are homeomorphic to a circle minus two points, so

$$Z_f^+(T) = 2(u-1) \frac{u^{-2} T^2}{1 - u^{-2} T^2} \frac{u^{-3} T^4}{1 - u^{-3} T^4} + (u-1) \frac{u^{-2} T^2}{1 - u^{-2} T^2} + (u-1) \frac{u^{-3} T^4}{1 - u^{-3} T^4}.$$

Note that in this case one has neither $Z_f(T) = (u-1) Z_f^+(T)$ nor $Z_f(T) = \frac{u-1}{2} Z_f^+(T)$, whereas it was the case in the preceding examples.

Remark 4.6. It would be convenient to have a Thom-Sebastiani formula to compute the zeta functions of the function $f * g$ defined by $(x, y) \rightarrow f(x) + g(y)$ from the ones of f and g , as it is the case in [15, 5, 21]. But it seems to be impossible to find in general such formulae with our zeta functions. However in the particular case of two positive (respectively negative) functions, one has the following formulae.

Proposition 4.7. *Let $f : \mathbb{R}^{d_1} \rightarrow \mathbb{R}$ and $g : \mathbb{R}^{d_2} \rightarrow \mathbb{R}$ be two positive or two negative functions. Let us put $Z_f(T) = \sum_{n \geq 1} a_n T^n$ and $Z_g(T) = \sum_{n \geq 1} b_n T^n$, and $A_n = 1 - \sum_{j=1}^n a_j$,*

$B_n = 1 - \sum_{j=1}^n b_j$. Then the naive zeta function of $f * g : \mathbb{R}^{d_1+d_2} \longrightarrow \mathbb{R}$ defined by $f * g(x, y) = f(x) + g(y)$ is $Z_{f*g}(T) = \sum_{n \geq 1} c_n T^n$ with

$$c_n = a_n B_n + A_n b_n + a_n b_n.$$

Example 4.8. (1) Let $h : \mathbb{R}^2 \longrightarrow \mathbb{R}$ be defined by $h(x, y) = x^2 + y^2$. Recall that $Z_f(T) = (u^2 - 1) \sum_{n \geq 1} \frac{T^{2n}}{u^{2n}}$ (cf example 4.3). Putting $f(x) = g(x) = x^2$, then $h = f * g$ and by example 4.1 we get that $a_{2n} = b_{2n} = \frac{u-1}{u^n}$ and $a_{2n+1} = b_{2n+1} = 0$, hence $A_{2n} = A_{2n+1} = \frac{1}{u^n}$. Then by proposition 4.7 we rederive $c_{2n} = \frac{u^2-1}{u^{2n}}$ and $c_{2n+1} = 0$.
(2) Consider f as in example 4.5.3. The odd coefficients are zero because f and g are positive, and it is easy to verify that

$$a_{2n} = \frac{u-1}{u^n}, A_{2n} = \frac{1}{u^n},$$

and

$$b_{4n} = \frac{u-1}{u^n}, b_{4n+2} = 0, B_{4n} = \frac{1}{u^n} = B_{4n+2}.$$

Therefore

$$c_{4n} = \frac{u^2-1}{u^{3n}}, c_{4n+2} = \frac{u-1}{u^{3n+1}},$$

which was not so clear on the Denef & Loeser formula.

Proof. Remark first that $u^{nd_1} A_n = \beta(\{\gamma \in \mathcal{L}_n : \text{ord}(f \circ \gamma) > n\})$. Actually

$$\mathcal{L}_n = \pi_{n,1}^{-1}(\chi_1) \sqcup \dots \sqcup \pi_{n,n}^{-1}(\chi_n) \sqcup \{\gamma \in \mathcal{L}_n : \text{ord}(f \circ \gamma) > n\},$$

hence by applying β one gets

$$u^{nd_1} = \sum_{j=1}^n a_j u^{nd_1} + \beta(\{\gamma \in \mathcal{L}_n : \text{ord}(f \circ \gamma) > n\})$$

and the result follows.

Now take $(\gamma_1, \gamma_2) \in \mathcal{L}_n(\mathbb{R}^{d_1}) \times \mathcal{L}_n(\mathbb{R}^{d_2}) = \mathcal{L}_n(\mathbb{R}^{d_1+d_2})$; then $\text{ord}(f \circ \gamma_1 + g \circ \gamma_2)$ is greater than n if and only if $\text{ord}(f \circ \gamma_1)$ and $\text{ord}(g \circ \gamma_2)$ are greater than n because f and g are of the same sign. Therefore we have to distinguish the cases $\text{ord}(f \circ \gamma_1) = n$ and $\text{ord}(g \circ \gamma_2) > n$, $\text{ord}(f \circ \gamma_1) > n$ and $\text{ord}(g \circ \gamma_2) = n$, and finally $\text{ord}(f \circ \gamma_1) = n$ and $\text{ord}(g \circ \gamma_2) = n$:

$$u^{n(d_1+d_2)} \beta(\chi_n(f * g)) = \beta(\chi_n(f)) u^{nd_2} B_n + u^{nd_1} A_n \beta(\chi_n(g)) + \beta(\chi_n(f)) \beta(\chi_n(g)).$$

□

4.2. Blow-Nash equivalence. Let $f, g : (\mathbb{R}^d, 0) \longrightarrow (\mathbb{R}, 0)$ be two germs of real analytic functions. They are said to be *blow-Nash* equivalent if there exist two modifications $\sigma_f : (M_f, \sigma_f^{-1}(0)) \longrightarrow (\mathbb{R}^d, 0)$ and $\sigma_g : (M_g, \sigma_g^{-1}(0)) \longrightarrow (\mathbb{R}^d, 0)$, that is proper birational maps which are isomorphisms over the complement of the zero locus of f and g , and a Nash-isomorphism (that is a semi-algebraic map which is an analytic isomorphism) Φ between $(M_f, \sigma_f^{-1}(0))$ and $(M_g, \sigma_g^{-1}(0))$ which preserves the multiplicities of the jacobian determinants of σ_f and σ_g along the components of the exceptional divisors, and which induces a homeomorphism $\phi : (\mathbb{R}^d, 0) \longrightarrow (\mathbb{R}^d, 0)$ such that $f = g \circ \phi$:

$$\begin{array}{ccc}
(M_f, \sigma_f^{-1}(0)) & \xrightarrow{\Phi} & (M_g, \sigma_g^{-1}(0)) \\
\sigma_f \downarrow & & \downarrow \sigma_g \\
(\mathbb{R}^d, 0) & \xrightarrow{\phi} & (\mathbb{R}^d, 0) \\
& \searrow f & \swarrow g \\
& (\mathbb{R}, 0) &
\end{array}$$

Blow-*Nash* equivalence is a particular case of blow-analytic equivalence, for which we add algebraic data. A difficult issue with this kind of equivalence relation is to find invariants, and actually we know just the Fukui invariants [12] and the Zeta functions of S. Koike and A. Parusiński [15], defined with the Euler characteristic with compact support.

Note that the naive zeta function generalizes the Fukui invariants which assign to f the possible orders of series $f \circ \gamma$ where γ is a real analytic arc at the origin. Indeed $\beta(\chi_n(f)) \neq 0$ as soon as $\chi_n(f) \neq \emptyset$ (recall remark 2.8), and thus the Fukui invariants are the nonzero exponents of $Z_f(T)$.

The following result is the main one of this section. Its proof is directly inspired from [15], theorem 4.5, and is a direct consequence of lemma 4.17 (see section 4.4 below).

Theorem 4.9. *The zeta functions $Z_f(T), Z_f^\pm(T)$ of a germ of analytic functions are invariants of blow-*Nash* equivalence.*

Proof. Let $f, g : (\mathbb{R}^d, 0) \longrightarrow (\mathbb{R}, 0)$ be two blow-*Nash* equivalent analytic function germs. By definition there exist modifications $\sigma_f : (M_f, \sigma_f^{-1}(0)) \longrightarrow (\mathbb{R}^d, 0)$ and $\sigma_g : (M_g, \sigma_g^{-1}(0)) \longrightarrow (\mathbb{R}^d, 0)$, and a Nash-isomorphism $\Phi : (M_f, \sigma_f^{-1}(0)) \longrightarrow (M_g, \sigma_g^{-1}(0))$. By [11, 2] one can assume that $f \circ \sigma_f$ and the jacobian determinant $\text{jac } \sigma_f$ (respectively $g \circ \sigma_g$ and the jacobian determinant $\text{jac } \sigma_g$) are normal crossing simultaneously, and thus the assumptions of lemma 4.17 are verified. Now It suffices to prove that the Denef & Loeser formulae for the zeta functions given by lemma 4.17 coincide. But β is invariant under Nash-isomorphism by theorem 3.3, and moreover ϕ preserves

- the multiplicities of $f \circ \sigma_f$ and $g \circ \sigma_g$ because it is an isomorphism
- the multiplicities of the jacobians of σ_f and σ_g along the components of the exceptional divisors by definition of blow-*Nash* equivalence.

Therefore the zeta functions of f and g coincide.

□

4.3. Application to Brieskorn polynomials. We apply our zeta functions to sketch the classification of two variables Brieskorn polynomials under blow-*Nash* equivalence, and to give examples in three variables. Brieskorn polynomials in two or three variables are polynomials of the type $\varepsilon_p x^p + \varepsilon_q y^q (+\varepsilon_r z^r)$, $p \leq q \leq r \in \mathbb{N}, \varepsilon_p, \varepsilon_q, \varepsilon_r \in \{\pm 1\}$.

Remark that if $p = 1$ then $\varepsilon_p x + \varepsilon_q y^q (+\varepsilon_r z^r)$ is *Nash*-isomorphic to x . Therefore we will restrict our attention to the case $p \geq 2$.

Actually the classification under blow-analytic equivalence has been done completely in the two variable case, and almost completely in the three variables case in [15] using zeta functions defined with Euler characteristic with compact support and the Fukui invariants(see [15], theorem 7.3; for the Fukui invariant, see [12]). There are just one case they can not

discuss, and the following example shows that we can discuss it under blow-*Nash* equivalence. However this it is not sufficient to conclude for blow-analytic equivalence.

Example 4.10. Let $f_{p,k}$ be the Brieskorn polynomial defined by $f_{p,k} = \pm(x^p + y^{kp} + z^{kp})$, p even, $k \in \mathbb{N}$. We are going to prove that for fixed p and different k , to such polynomials are not blow-*Nash* equivalent.

In order to do this we calculate directly the zeta function of $f_{p,k}$. For $n \in \mathbb{N}$ we compute $\beta(\chi_n)$. First it is clear that $\chi_n = \emptyset$ when n is not a multiple of p . If n is a multiple of p , write $n = p(mk + r)$ with $mk + r$ the euclidean division of $\frac{n}{p}$ by k . If $\gamma \in \mathcal{L}_n$, put $\gamma = (a_1t + \dots + a_nt^n, b_1t + \dots + b_nt^n, c_1t + \dots + c_nt^n)$.

Then if $r \neq 0$, the first non zero term of $f \circ \gamma$ is given by the first component of γ , hence

$$\chi_n = \{\gamma | a_{mk+r} \neq 0, a_1 = \dots = a_{mk+r-1} = b_1 = \dots = b_m = c_1 = \dots = c_m = 0\}.$$

In the case where $r = 0$, the three components of γ play a part, and

$$\chi_n = \{\gamma | (a_{mk}, b_m, c_m) \neq (0, 0, 0), a_1 = \dots = a_{mk-1} = b_1 = \dots = b_{m-1} = c_1 = \dots = c_{m-1} = 0\}.$$

Therefore

$$\chi_n \simeq \begin{cases} \mathbb{R}^* \times \mathbb{R}^{(p-1)(mk+r)} \times (\mathbb{R}^{p(mk+r)-m})^2 & \text{if } r \neq 0 \\ (\mathbb{R}^3)^* \times \mathbb{R}^{(p-1)mk} \times (\mathbb{R}^{pmk-m})^2 & \text{if } r = 0 \end{cases}$$

hence the coefficient of T^n is

$$\beta(\chi_n)u^{-3n} = \begin{cases} (u-1)u^{-(mk+r)-2m} & \text{if } n = p(mk+r), 0 < r < k \\ (u^3-1)u^{-mk-2m} & \text{if } n = pmk \\ 0 & \text{otherwise.} \end{cases}$$

Therefore the zeta function of $f_{p,k}$ looks like

$$Z_{f_{p,k}} = (u-1)(u^{-1}T^p + u^{-2}T^{2p} + \dots + u^{-(k-1)}T^{(k-1)p}) + (u^3-1)u^{-k-2}T^{kp} \\ + (u-1)(u^{-(k+3)}T^{(k+1)p} + u^{-(k+4)}T^{(k+2)p} + \dots + u^{-(2k+1)}T^{(2k-1)p}) + (u^3-1)u^{-2(k-2)}T^{2kp} + \dots$$

Now it suffices to note that, for p fixed and $k < k'$ the pk -coefficient of $Z_{f_{p,k}}$ is $(u^3-1)u^{-k-2}$ whereas the one of $Z_{f_{p,k'}}$ is $(u-1)u^{-k}$.

Remark 4.11. The case of two variables Brieskorn polynomials have been dealt with in [15], using their zeta functions and the Fukui invariants. Actually the only case where the equivalence class of Brieskorn polynomials of two variables can not be distinguished using only their zeta functions, and which requires the use of the Fukui invariants, is the following: $f_k(x, y) = \pm(x^k + y^k)$, $k \geq 2$ even. Remark that we have seen in example 4.3 that for $k \neq k'$ the zeta function Z_{f_k} and $Z_{f_{k'}}$ are different, therefore our zeta function distinguishes this case, for blow-*Nash* equivalence.


Actually, one can say more. Indeed, for two variables Brieskorn polynomials the naive zeta function determines the coefficients p and q .

Proposition 4.12. *Let $g = \pm x^p \pm y^q$ be a two variable Brieskorn polynomial. Then p and q are uniquely determined by the naive zeta function $Z_g(T) = \sum_{n \geq 1} g_n T^n$. More precisely*

$$p = \min\{n : g_n \neq 0\}$$

and if $l = \min\{n : g_n \neq a_n u^n\}$, where $\sum_{n \geq 1} a_n T^n$ denote the naive zeta function of $\pm x^p$, then

FIGURE 4. Resolution tree of $x^3 \pm y^4$.



$$q = \begin{cases} l-1 & \text{if } p \text{ is odd, } p|l-1 \text{ and } g_l \neq (u-1)u^k \text{ for some } k \in \mathbb{N} \\ l & \text{otherwise.} \end{cases}$$

Proof. The characterization of p is clear. Now, if $p \nmid q$, then $q = l$ and $g_q = (u-1)u^k$ for some $k \in \mathbb{N}$.

If $q = kp$ for some $k \in \mathbb{N}$, then $g_{kp} = \beta(\{\pm a^p \pm b^{kp}\})u^{2kp-k-1}$ and thus

$$g_{kp} = a_{kp}u^{kp} \iff \beta(\{\pm a^p \pm b^{kp} \neq 0\}) = u(u-1) \iff p \text{ odd.}$$

In that case $g_{kp+1} = (u-1)^2u^{2kp-k-2}$, so $l = kp + 1$.

Therefore if p is even $q = l$, and if p is odd either $q = l$ or $q = l-1$. More precisely if $g_l = (u-1)u^k$ then $p \nmid q$ and $q = l$, whereas if $g_l = (u-1)^2u^k$, then $p|l-1$ and $q = l-1$. $\square$

To find the signs in front of x^p and y^q , the naive zeta function is not sufficient as shown by the following example.

Example 4.13. Let $f_{\pm} : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}, 0)$ be defined by $f_{\pm}(x, y) = x^3 \pm y^4$. One can solve the singularities of $f_{\pm}$ by a succession of four blowing-ups. The resolution tree of this modification σ is drawn in figure 4, where $Z_{\pm}$ denotes the strict transform of $f_{\pm}$, and $E(N, \nu)$ denote an irreducible component of the exceptional divisor such that $\text{mult}_E f_{\pm} \circ \sigma = N$ and $1 + \text{mult}_E \det \text{jac } \sigma = \nu$. By Denef & Loeser formula $Z_{f_+} = Z_{f_-}$, but f_+ and f_- are not blow-Nash equivalent (they are not even blow-analytically equivalent, see [15], theorem 6.1 for example).

Actually the zeta functions with signs enable to discuss the signs for two variables Brieskorn polynomials except in one case. The following proposition details the possibilities. Note that $(x, y) \rightarrow (\pm x, \pm y)$ give an action on the blow-Nash classes, hence when the coefficient p or q is odd, the corresponding sign can not be determined. By convention, in the case $p = q$ and the sign are opposite, we consider $x^p - y^p$ rather than $-x^p + y^p$.

Proposition 4.14. Let $Z_g(T) = \sum_{n \geq 1} g_n T^n$ be the naive motivic zeta function of $\varepsilon_p x^p + \varepsilon_q y^q$, with $\varepsilon_p, \varepsilon_q \in \{\pm 1\}$. If p is even

$$\varepsilon_p = \begin{cases} +1 & \text{if } g_p^+ \neq 0 \\ -1 & \text{otherwise,} \end{cases}$$

and if q is even, but not multiple of an odd p , then

$$\varepsilon_q = \begin{cases} +1 & \text{if } g_p^- = 0 \\ -1 & \text{otherwise.} \end{cases}$$

Remark 4.15. In the case p odd and $q = kp$ with k even, we do not know if $x^p + y^{kp}$ and $x^p - y^{kp}$ are blow-*Nash* equivalent or not. They are blow-analytically equivalent by an argument of integration of vector field, but this method no longer holds in the *Nash* setting.

4.4. Motivic integration and the proof of Denef & Loeser formula. The proof of Denef & Loeser formula, which is a simplification of the one of [8], theorem 2.2.1 to our setting, uses the theory of motivic integration on arc spaces for real algebraic varieties (for motivic integration, we refer to [5, 6, 7, 8, 16, 21]). In particular we will use the change of variables formula, which is a fundamental result for this theory ([16], and [6], lemma 3.3). We will recall these notions, without proofs.

Let $\sigma : (M, \sigma^{-1}(0)) \longrightarrow (\mathbb{R}^d, 0)$ be a real modification. Let

$$\mathcal{L}(M, \sigma^{-1}(0)) = \{\gamma : (\mathbb{R}, 0) \longrightarrow (M, \sigma^{-1}(0)) \mid \gamma \text{ is analytic}\}$$

be the arc space associated to $(M, \sigma^{-1}(0))$. Denote by $\pi_n : \mathcal{L} \longrightarrow \mathcal{L}_n$ the natural truncation morphisms, for $n \in \mathbb{N}$, where $\mathcal{L}$ denote either $\mathcal{L}(M, \sigma^{-1}(0))$ or $\mathcal{L}(\mathbb{R}^d, 0)$. A subset $A \subset \mathcal{L}$ is called stable if $A = \pi_n^{-1}(C)$ for $C \subset \mathcal{L}_n$ constructible and some $n \geq 0$. Then we can define the measure of such a stable set A in $\mathbb{Z}[u, u^{-1}]$ by

$$\beta(A) = u^{-(n+1)d} \beta(\pi_n(A))$$

for n large enough ($\beta(\pi_n(A))$ is well defined for Zariski constructible real algebraic varieties are arc-symmetric sets). Indeed, $\beta(A)$ does not depend on n because the natural projections $\mathcal{L}_{n+1} \longrightarrow \mathcal{L}_n$ are locally trivial fibrations with fiber $\mathbb{R}^d$.

Let $\theta : A \longrightarrow \mathbb{Z}[u, u^{-1}]$ be a map with finite image and whose fibers are stable sets. Then the integral of θ over A with respect to β is defined by the formula

$$\int_A \theta d\beta := \sum_{c \in \mathbb{Z}[u, u^{-1}]} c \beta(\theta^{-1}(c)).$$

We can state now the Kontsevich change of variables formula.

Proposition 4.16. ([16, 6]) *Let $A \subset \mathcal{L}(\mathbb{R}^d, 0)$ be stable, and suppose that $\text{ord}_t \det(\text{jac } \sigma)$ is bounded on $\sigma^{-1}(A)$. Then*

$$\beta(A) = \int_{\sigma^{-1}(A)} u^{-\text{ord}_t \det(\text{jac } \sigma)} d\beta.$$

Before giving the proof of Denef & Loeser formula, let us give some notations. The modification σ induces applications σ_* (respectively $\sigma_{*,n}$) between $\mathcal{L}(M, \sigma^{-1}(0))$ and $\mathcal{L}(\mathbb{R}^d, 0)$ (respectively $\mathcal{L}_n(M, \sigma^{-1}(0))$ and $\mathcal{L}_n(\mathbb{R}^d, 0)$). We put $\mathcal{Z}_n(f) = \pi_n^{-1}(\chi_n)$ and $\mathcal{Z}_n(f \circ \sigma) = \sigma_*^{-1}(\mathcal{Z}_n(f))$. Finally for $e \geq 1$, put $\Delta_e = \{\gamma \in \mathcal{L}(M, \sigma^{-1}(0)) \mid \text{ord}_t \det(\text{jac } \sigma)(\gamma(t)) = e\}$, and $\mathcal{Z}_{n,e}(f \circ \sigma) = \mathcal{Z}_n(f \circ \sigma) \cap \Delta_e$.

Let us state a preliminary lemma.

Lemma 4.17. *Let $\sigma : (M, \sigma^{-1}(0)) \longrightarrow (\mathbb{R}^d, 0)$ be a modification of $\mathbb{R}^d$ such that $f \circ \sigma$ and the jacobian determinant $\text{jac } \sigma$ are normal crossings simultaneously, and assume moreover that σ is an isomorphism over the complement of the zero locus of f .*

Let $(f \circ \sigma)^{-1}(0) = \cup_{j \in J} E_j$ be the decomposition into irreducible components of $(f \circ \sigma)^{-1}(0)$, and assume that $\sigma^{-1}(0) = \cup_{k \in K} E_k$ for some $K \subset J$.

Put $N_i = \text{mult}_{E_i} f \circ \sigma$ and $\nu_i = 1 + \text{mult}_{E_i} \text{jac } \sigma$, and for $I \subset J$ denote by E_I^0 the set $(\cap_{i \in I} E_i) \setminus (\cup_{j \in J \setminus I} E_j)$. Then there exists $c \in \mathbb{N}$ such that

$$Z_f(T) = u^d \sum_{n \geq 1} T^n \sum_{e \leq cn} u^{-e} \sum_{I \neq \emptyset} \beta(\{\gamma \in \mathcal{L}_n(M, E_I^0) \cap \pi_n(\Delta_e) : f \circ \sigma \circ \gamma(t) = bt^n + \dots, b \neq 0\})$$

and

$$Z_f^\pm(T) = u^d \sum_{n \geq 1} T^n \sum_{e \leq cn} u^{-e} \sum_{I \neq \emptyset} \beta(\{\gamma \in \mathcal{L}_n(M, E_I^0) \cap \pi_n(\Delta_e) : f \circ \sigma \circ \gamma(t) = \pm t^n + \dots\}).$$

Proof. Let us prove the lemma for $Z_f(T)$, the argument is the same for $Z_f^\pm$.

For $n \geq 1$, $\mathcal{Z}_n(f)$ is stable, so $\beta(\mathcal{Z}_n(f))$ is defined and equals $u^{-(n+1)d} \beta(\chi_n)$, hence $Z_f(T) = u^d \sum_{n \geq 1} \beta(\mathcal{Z}_n(f)) T^n$.

Moreover $\mathcal{Z}_n(f \circ \sigma)$ equals the disjoint union $\cup_{e \geq 1} \mathcal{Z}_{n,e}(f \circ \sigma)$. Actually this union is finite: take $\gamma \in \mathcal{Z}_n(f \circ \sigma)$; there exists $I \subset J$ such that $\pi_0(\gamma) \in E_I^0$. Then in a neighbourhood of $\gamma(0)$ one can choose coordinates such that $f \circ \sigma = \text{unit} \prod_{i \in I} y_i^{N_i}$ and $\det(\text{jac } \sigma) = \text{unit} \prod_{i \in I} y_i^{\nu_i - 1}$ where by *unit* we denote a non-vanishing function. Let us write $\gamma = (\gamma_1, \dots, \gamma_d)$, and $k_i = \text{ord}_t \gamma_i$, for $i = 1, \dots, d$. Then $\text{ord}_t f \circ \sigma(\gamma(t)) = \sum_{i=1}^d N_i k_i \geq n$ and therefore

$$\text{ord}_t \det(\text{jac } \sigma)(\gamma(t)) = \sum_{i=1}^d (\nu_i - 1) k_i \geq \max_i \left(\frac{\nu_i - 1}{N_i} \right) \sum_{i=1}^d N_i k_i \geq \max_i \left(\frac{\nu_i - 1}{N_i} \right) n.$$

Putting $c = \max_i \left(\frac{\nu_i - 1}{N_i} \right)$, we have shown thus that $\cup_{e \geq 1} \mathcal{Z}_{n,e}(f \circ \sigma) = \cup_{e \leq cn} \mathcal{Z}_{n,e}(f \circ \sigma)$ which is a finite union.

Now Kontsevich change of variables formula induces that

$$\beta(\mathcal{Z}_n(f)) = \sum_{e \leq cn} u^{-e} \beta(\mathcal{Z}_{n,e}(f \circ \sigma)),$$

and then

$$Z_f(T) = u^d \sum_{n \geq 1} T^n \sum_{e \leq cn} u^{-e} \beta(\mathcal{Z}_{n,e}(f \circ \sigma)).$$

We are going to compute $\beta(\mathcal{Z}_{n,e}(f \circ \sigma))$ using the fact that $\mathcal{Z}_{n,e}(f \circ \sigma)$ equals the disjoint union

$$\sqcup_{I \neq \emptyset} \mathcal{Z}_{n,e}(f \circ \sigma) \cap \pi_0^{-1}(E_I^0 \cap \sigma^{-1}(0)),$$

thus

$$\beta(\mathcal{Z}_{n,e}(f \circ \sigma)) = \sum_{I \neq \emptyset} \beta(\mathcal{Z}_{n,e}(f \circ \sigma) \cap \pi_0^{-1}(E_I^0 \cap \sigma^{-1}(0))).$$

Choose $I \neq \emptyset$. Then $\pi_n(\mathcal{Z}_{n,e}(f \circ \sigma) \cap \pi_0^{-1}(E_I^0 \cap \sigma^{-1}(0)))$ is just the set

$$\{\gamma \in \mathcal{L}_n(M, \sigma^{-1}(0)) \mid \gamma(0) \in E_I^0 \cap \sigma^{-1}(0), \text{ord}_t f \circ \sigma(\gamma(t)) = n, \text{ord}_t \det(\text{jac } \sigma)(\gamma(t)) = e\}.$$

The results follows directly from the additivity of β .

□

The proofs of propositions 4.2 and 4.4 just consist of computing the value of β on the sets which appear in the formulae of lemma 4.17. Let us first prove proposition 4.2.

Proof of proposition 4.2. Take $\gamma \in \pi_n(\mathcal{Z}_{n,e}(f \circ \sigma) \cap \pi_0^{-1}(E_I^0 \cap \sigma^{-1}(0)))$. On a neighbourhood of $\gamma(0)$ one can choose coordinates such that $f \circ \sigma = \text{unit} \prod_{i \in I} y_i^{N_i}$ and moreover $\det(\text{jac } \sigma) = \text{unit} \prod_{i \in I} y_i^{\nu_i - 1}$, hence $\pi_n(\mathcal{Z}_{n,e}(f \circ \sigma) \cap \pi_0^{-1}(E_I^0 \cap \sigma^{-1}(0)))$ is isomorphic to

$$\{\gamma \in \mathcal{L}_n(M, \sigma^{-1}(0)) \mid \gamma(0) \in E_I^0 \cap \sigma^{-1}(0), \sum_{i \in I} k_i N_i = n, \sum_{i \in I} k_i (\nu_i - 1) = e\},$$

where $k_i = \text{ord}_t \gamma_i$ for $i \in I$. As a consequence

$$\pi_n(\mathcal{Z}_{n,e}(f \circ \sigma) \cap \pi_0^{-1}(E_I^0 \cap \sigma^{-1}(0))) \simeq \sqcup_{k \in A(n,e)} (E_I^0 \cap \sigma^{-1}(0)) \times (\mathbb{R}^*)^{|I|} \left(\prod_{i \in I} \mathbb{R}^{n-k_i} \right) \times (\mathbb{R}^n)^{d-|I|}$$

where $A(n,e)$ is the subset of $k \in \mathbb{N}^d$ such that $\sum_{i=1}^d N_i k_i = n$ and $\sum_{i=1}^d (\nu_i - 1) k_i = e$.

By taking the image by β we obtain

$$\beta\left(\pi_n(\mathcal{Z}_{n,e}(f \circ \sigma) \cap \pi_0^{-1}(E_I^0 \cap \sigma^{-1}(0)))\right) = \beta(E_I^0 \cap \sigma^{-1}(0)) (u-1)^{|I|} u^{nd - \sum_{i=1}^d k_i},$$

hence

$$Z_f(T) = \sum_{I \neq \emptyset} (u-1)^{|I|} \beta(E_I^0 \cap \sigma^{-1}(0)) \sum_{n \geq 1} \sum_{e \leq cn} \sum_{k \in A(n,e)} u^{-e - \sum_{i=1}^d k_i} T^n.$$

Remark that $\{k \in A(n,e) \mid n \geq 1, e \leq cn\}$ is in bijection with $\mathbb{N}^{|I|}$, therefore

$$\sum_{n \geq 1} \sum_{e \leq cn} \sum_{k \in A(n,e)} u^{-e - \sum_{i=1}^d k_i} T^n = \sum_k \prod_{i \in I} (u^{-\nu_i} T^{N_i})^{k_i} = \prod_{i \in I} \frac{u^{-\nu_i} T^{N_i}}{1 - u^{-\nu_i} T^{N_i}}.$$

Finally

$$Z_f(T) = \sum_{I \neq \emptyset} (u-1)^{|I|} \beta(E_I^0 \cap \sigma^{-1}(0)) \prod_{i \in I} \frac{u^{-\nu_i} T^{N_i}}{1 - u^{-\nu_i} T^{N_i}},$$

which is the result. $\square$

The proof of proposition 4.4 is a little bit more complicated due to the fact that we have to introduce a covering $\widetilde{E_I^{0,\pm}}$ of $E_I^{0,\pm}$ in order to compute $Z_f^\pm(T)$. Recall that if U is an affine open subset of M such that $f \circ \sigma = u \prod_{i \in I} y_i^{N_i}$ on U , with u a unit, then by $R_U^\pm$ we mean $R_U^\pm = \{(x, t) \in (E_I^0 \cap U) \times \mathbb{R} : t^m = \pm \frac{1}{u(x)}\}$, where $m = \gcd(N_i)$. Now $\widetilde{E_I^{0,\pm}}$ is the gluing of the $R_U^\pm$ along the $E_I^0 \cap U$.

Proof of proposition 4.4. Let U be an affine open subset of M such that $f \circ \sigma = u \prod_{i \in I} y_i^{N_i}$ on U , with u a unit. What we have to compute is the value of β on

$$W^\pm = \{(x, y) \in (E_I^0 \cap U) \times (\mathbb{R}^*)^{|I|} : u(x) \prod_{i \in I} y_i^{N_i} = \pm 1\}.$$

Denote by m the greatest common divisor of the N_i , $i \in I$, and choose n_i , $i \in I$ such that $\sum_{i \in I} n_i N_i = m$. Assume that $I = \{1, \dots, s\}$. Remark that $W^\pm$ is isomorphic to

$$W'^{\pm} = \{(x, y, t) \in (E_I^0 \cap U) \times (\mathbb{R}^*)^{|I|} \times \mathbb{R}^* : t^m = \frac{\pm 1}{u(x)}, \prod_{i \in I} y_i^{N_i/m} = 1\},$$

by $W'^{\pm} \longrightarrow W^{\pm}, (x, y, t) \longrightarrow (x, t^{n_1}y_1, \dots, t^{n_s}y_s)$. The inverse is the morphism given by $(x, y) \longrightarrow (x, \prod_{i \in I} y_i^{N_i/m}, (\prod_{i \in I} y_i^{N_i/m})^{-n_1}y_1, \dots, (\prod_{i \in I} y_i^{N_i/m})^{-n_s}y_s)$.

Now it is easier to compute $\beta(W'^{\pm})$ for $W'^{\pm} \simeq R_U^{\pm} \times (\mathbb{R}^*)^{|I|-1}$. This last isomorphism comes from the fact that at least one $\frac{N_i}{m}$ is odd. Therefore $\beta(W^{\pm}) = (u-1)^{|I|-1} \beta(R_U^{\pm})$, and the same computation as in the naive case gives the formula.

□

REFERENCES

- [1] D. Abramovich, K. Karu, K. Matsuki and J. Włodarczyk, *Torification and factorization of birational maps*, J. Amer. Math. Soc. 15, no. 3, 531–572, (2002)
- [2] E. Bierstone, P.D. Milman, *Canonical desingularization in characteristic zero by blowing up the maximum strata of a local invariant*, Invent. math. 128, 207–302 (1997)
- [3] F. Bittner, *The universal Euler characteristic for varieties of characteristic zero*, Compositio Math. (to appear)
- [4] J. Bochnak, M. Coste, M.-F. Roy, *Real Algebraic Geometry*, Springer-Verlag, Berlin 1998
- [5] J. Denef, F. Loeser, *Geometry on arc spaces of algebraic varieties* European Congress of Math. (Barcelona, July 10–14, 2000) 1 (2001), 325–348
- [6] J. Denef, F. Loeser, *Germes of arcs on singular algebraic varieties and motivic integration*, Invent. math. 135 (1999), 201–232
- [7] J. Denef, F. Loeser, *Lefschetz numbers of iterates of the monodromy and truncated arcs*,
- [8] J. Denef, F. Loeser, *Motivic Igusa zeta functions*, J. Algebraic Geom. 7 (1998), 3, 505–537
- [9] T. Fukui, *Seeking invariants for blow-analytic equivalence*, Compositio Math. 105 (1997), 95–107
- [10] T. Fukui, S. Koike, T.-C. Kuo, *Blow-analytic equisingularities, properties, problems and progress*, Real Analytic and Algebraic Singularities (T. Fukuda, T. Fukui, S. Izumiya and S. Koike, ed), Pitman Research Notes in Mathematics Series, 381 (1998), pp. 8–29
- [11] H. Hironaka, *Resolution of singularities of an algebraic variety over a field of characteristic zero*, Ann. Math. 79, 109–326 (1964)
- [12] S. Izumi, S. Koike, T.-C. Kuo, *Computations and stability of the Fukui Invariant*, Compositio Math. 130 (2002), 49–73
- [13] S. Koike, *Modified Nash triviality theorem for a family of zero-sets of weighted homogeneous polynomial mappings*, J. Math. Soc. Japan 49 (1997), 617–631
- [14] S. Koike, *Nash trivial simultaneous resolution for a family of zero-sets of Nash mappings*, Math. Zeitschrift 234 (2000), 313–338
- [15] S. Koike, A. Parusiński, *Motivic-type invariants of blow-analytic equivalence*, Ann. Inst. Fourier (to appear)
- [16] M. Kontsevich, *Lecture at Orsay* (December 7, 1995)
- [17] T.-C. Kuo, *On classification of real singularities*, Invent. Math. 82 (1985), 257–262
- [18] K. Kurdyka, *Ensembles semi-algébriques symétriques par arcs*, Math. Ann. 282, 445–462 (1988)
- [19] K. Kurdyka, *Injective endomorphisms of real algebraic sets are surjective*, Math. Ann. 313, 69–82 (1999)
- [20] S. Lojasiewicz, *Introduction to complex analytic geometry*, Birkhauser, 1991
- [21] E. Looijenga, *Motivic Measures*, in Séminaire Bourbaki, exposé 874, Mars 2000
- [22] C. McCrory, A. Parusiński, *Virtual Betti numbers of real algebraic varieties*, C. R. Acad. Sci. Paris Sr. I Math. (to appear)
- [23] A. Parusiński, *Constructible categories and the Borel-Kurdyka theorem*, University of Angers preprint no. 148, April 2002
- [24] R. Quarez, *Espace des germes d’arcs réels et série de Poincaré d’un ensemble semi-algébrique*, Ann. Inst. Fourier (Grenoble) 51 (2001), no. 1, 43–68.
- [25] B. Totaro, *Topology of singular algebraic varieties*, Proc. Int. Cong. Math. Beijing, vol. 1 (2002), 533–541
- [26] J. Włodarczyk, *Toroidal varieties and the weak factorization theorem*, math.AG/9904076

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ D’ANGERS, 2 BD LAVOISIER, 49045 ANGERS CEDEX, FRANCE

E-mail address: fichou@tonton.univ-angers.fr